

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/475, A61K 37/02		A1	(11) International Publication Number: WO 96/27007 (43) International Publication Date: 6 September 1996 (06.09.96)
(21) International Application Number: PCT/AU96/00094 (22) International Filing Date: 22 February 1996 (22.02.96) (30) Priority Data: PN 1457 2 March 1995 (02.03.95) AU PN 6647 20 November 1995 (20.11.95) AU PN 7274 22 December 1995 (22.12.95) AU (71) Applicant (for all designated States except US): AMRAD OPERATIONS PTY. LTD. [AU/AU]; 17-27 Cotham Road, Kew, VIC 3101 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only): HAYWARD, Nicholas, Kim [AU/AU]; 13 Prince Street, Paddington, QLD 4064 (AU). WEBER, Gunther [SE/AU]; 17-27 Cotham Road, Kew, VIC 3101 (AU). GRIMMOND, Sean [AU/AU]; 210 Swann Road, Taringa, QLD 4068 (AU). NORDEN-SKJOLD, Magnus [SE/SE]; Koltrastvagen 19, S-161 37 Stockholm (SE). LARSSON, Catharina [SE/SE]; Dannemoragatan 10m 3tr, S-113 44 Stockholm (SE). (74) Agents: HUGHES, E., John, L. et al.; Davies Collison Cave, 1 Little Collins Street, Melbourne, VIC 3000 (AU).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: A NOVEL GROWTH FACTOR AND A GENETIC SEQUENCE ENCODING SAME			
(57) Abstract <p>The present invention relates generally to an isolated molecule having vascular endothelial growth factor-like properties and to a genetic sequence encoding same. The molecule will be useful in the development of a range of therapeutics and diagnostics useful in the treatment, prophylaxis and/or diagnosis of conditions requiring enhanced or diminished vasculature and/or vascular permeability. The molecule of the present invention is also a useful effector of primary and central neurons and is capable of inducing astroglial proliferation.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

A NOVEL GROWTH FACTOR AND A GENETIC SEQUENCE
ENCODING SAME

5 The present invention relates generally to an isolated molecule having vascular endothelial growth factor-like properties and to a genetic sequence encoding same. The molecule will be useful in the development of a range of therapeutics and diagnostics useful in the treatment, prophylaxis and/or diagnosis of conditions requiring enhanced or diminished vasculature and/or vascular permeability. The molecule of the present
10 invention is also a useful effector of primary and central neurons and is capable of inducing astroglial proliferation.

Bibliographic details of the publications referred to by author in this specification are collected at the end of the description. Sequence Identity Numbers (SEQ ID NOs.) for
15 the nucleotide and amino acid sequences referred to in the specification are defined following the bibliography.

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to
20 imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

Vascular endothelial growth factor (hereinafter referred to as "VEGF"), also known as vasoactive permeability factor, is a secreted, covalently linked homodimeric glycoprotein
25 that specifically activates endothelial tissues (Senger *et al.*, 1993). A range of functions have been attributed to VEGF such as its involvement in normal angiogenesis including formation of the corpus luteum (Yan *et al.*, 1993) and placental development (Sharkey *et al.*, 1993), regulation of vascular permeability (Senger *et al.*, 1993), inflammatory angiogenesis (Sunderkotter *et al.*, 1994) and autotransplantation (Dissen *et al.*, 1994) and
30 human diseases such as tumour promoting angiogenesis (Folkman & Shing, 1992), rheumatoid arthritis (Koch *et al.*, 1994) and diabetes related retinopathy (Folkman & Shing, 1992).

- 2 -

VEGF is, therefore, an important molecule making it a potentially valuable target for research into therapeutics, prophylactics and diagnostic agents based on VEGF or its activities. There is also a need to identify homologues or otherwise related molecules for use as an alternative to VEGF or in conjunction with VEGF.

5

In work leading up to the present invention, the inventors sought the multiple endocrine neoplasia type I susceptibility gene (MEN1). Surprisingly, the inventors discovered that a genetic sequence excluded as a candidate for the MEN1 gene was nevertheless a new growth factor having some similarity to VEGF. Furthermore, the growth factor of the present invention is an effector molecule for primary and central neurons.

10

Accordingly, one aspect of the present invention comprises a biologically isolated proteinaceous molecule comprising a sequence of amino acids which:

- (i) is at least about 15% similar to the amino acid sequence set forth in SEQ ID NO:2; and
- (ii) is at least 5% dissimilar to the amino acid sequence set forth in SEQ ID NO:2.

15

Another aspect of the present invention provides a biologically isolated proteinaceous molecule having the following characteristics:

- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to all or part of the amino acid sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one property in common with VEGF.

20

A related aspect of the present invention contemplates a biologically isolated proteinaceous molecule having the following characteristics:

- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the amino acid sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one of the following properties:
 - (a) ability to induce proliferation of vascular endothelial cells;
 - (b) ability to interact with *flt-1/flk-1* family of receptors;

25
30

- 3 -

- (c) ability to induce cell migration, cell survival and/or an increase in intracellular levels of alkaline phosphatase.

By "biologically isolated" is meant that the molecule has undergone at least one step of purification from a biological source. Preferably, the molecule is also biologically pure meaning that a composition comprises at least about 20%, more preferably at least about 40%, still more preferably at least about 65%, even still more preferably at least about 80-90% or greater of the molecule as determined by weight, activity or other convenient means, relative to other compounds in the composition. Most preferably, the molecule is sequencably pure.

Another preferred aspect of the present invention provides the molecule in recombinant form.

According to this aspect of the present invention, there is provided a recombinant molecule comprising a sequence of amino acids which:

- (i) is at least about 15% similar to the amino acid sequence set forth in SEQ ID NO:2; and
- (ii) is at least 5% dissimilar to the amino acid sequence set forth in SEQ ID NO:2.

A related aspect of the present invention is directed to a recombinant molecule having the following characteristics:

- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to all or part of the amino acid sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one property in common with VEGF.

A further related aspect of the present invention contemplates a recombinant molecule having the following characteristics:

- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the amino acid sequence set forth in SEQ ID NO:2;

- 4 -

(ii) exhibits at least one of the following properties:

- (a) ability to induce proliferation of vascular endothelial cells;
 - (b) ability to interact with *flt-1/flk-1* family of receptors;
 - (c) ability to induce cell migration, cell survival and/or an increase in
- 5 intracellular levels of alkaline phosphatase.

The present invention also contemplates genomic or partial genome clones encoding a proteinaceous molecule having at least about 15% amino acid similarity but at least about 5% dissimilarity to SEQ ID NO:1.

10

The amino acid sequence set forth in SEQ ID NO:2 corresponds to human VEGF (referred to herein as "VEGF₁₆₅"). Accordingly, the molecule of the present invention is VEGF-like or is a homologue of VEGF but comprises an amino acid sequence which is similar but non-identical to the amino sequence of VEGF. Although the present

15 invention is exemplified using a human VEGF-like molecule, this is done with the understanding that the instant invention contemplates the homologous molecule and encoding sequence from other mammals such as livestock animals (e.g. sheep, pigs, horses and cows), companion animals (e.g. dogs and cats) and laboratory test animals (e.g. mice, rats, rabbits and guinea pigs) as well as non-mammals such as birds (e.g.

20 poultry birds), fish and reptiles. In a most preferred embodiment, the VEGF-like molecule is of human origin and encoded by a gene located at chromosome 11q13. The present invention extends, therefore, to the genomic sequence or part thereof encoding the subject VEGF-like molecule.

25 Preferably, the percentage similarity is at least about 30%, more preferably at least about 40%, still more preferably at least about 50%, still even more preferably at least about 60-70%, yet even more preferably at least about 80-95% to all or part of the amino acid sequence set forth in SEQ ID NO:2.

30 In a particularly preferred embodiment, the VEGF-like molecule of the present invention comprises a sequence of amino acids as set forth in SEQ ID NO:4 or is a part, fragment, derivative or analogue thereof. Particularly preferred similarities include about 19-20%,

- 5 -

and 29-30%. Reference herein to derivatives also includes splice variants. Accordingly, the present invention extends to splice variants of SOM175. Examples of splice variants contemplated by the present invention include but are not limited to variants with an amino acid sequence substantially as set forth in at least one of SEQ ID NO:6, SEQ ID
5 NO:8 and/or SEQ ID NO:10 or mutants or derivatives or further splice variants thereof.

Another embodiment provides a recombinant molecule having the following characteristics:

- 10 (i) an amino acid sequence substantially as set forth in SEQ ID NO:4 or having at least about 15% similarity to all or part thereof provided that said amino acid sequence is at least about 5% dissimilar to all or part of the amino acid sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one biological property in common with VEGF.

15 Another embodiment provides a recombinant molecule having the following characteristics:

- 20 (i) an amino acid sequence substantially as set forth in SEQ ID NO:6 or having at least about 15% similarity to all or part thereof provided that said amino acid sequence is at least about 5% dissimilar to all or part of the amino acid sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one biological property in common with VEGF.

Another embodiment provides a recombinant molecule having the following characteristics:

- 25 (i) an amino acid sequence substantially as set forth in SEQ ID NO:8 or having at least about 15% similarity to all or part thereof provided that said amino acid sequence is at least about 5% dissimilar to all or part of the amino acid sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one biological property in common with VEGF.

30

Another embodiment provides a recombinant molecule having the following characteristics:

- 6 -

- (i) an amino acid sequence substantially as set forth in SEQ ID NO:10 or having at least about 15% similarity to all or part thereof provided that said amino acid sequence is at least about 5% dissimilar to all or part of the amino acid sequence set forth in SEQ ID NO:2;
- 5 (ii) exhibits at least one biological property in common with VEGF.

Such properties of VEGF include at least one of:

- (a) ability to induce proliferation of vascular endothelial cells;
- (b) an ability to interact with *flt-1/flk-1* family of receptors;
- 10 (c) an ability to induce cell migration, cell survival and/or an increase in intracellular levels of alkaline phosphatase.

In accordance with the present invention, a preferred similarity is at least about 40%, more preferably at least about 50% and even more preferably at least about 65% similarity.

15

Still a further aspect of the present invention contemplates a peptide fragment corresponding to a portion of the amino acid sequence set forth in SEQ ID NO:4 or a splice variant thereof such as set forth in SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:10 or a chemical equivalent thereof. The biologically isolated or recombinant molecule of the present invention may be naturally glycosylated or may comprise an altered glycosylation pattern depending on the cells from which it is isolated or synthesised. For example, if produced by recombinant means in prokaryotic organisms, the molecule would be non-glycosylated. The molecule may be a full length, naturally occurring form or may be a truncated or otherwise derivatised form.

20

25

Yet another aspect of the present invention is directed to a nucleic acid molecule encoding the VEGF-like molecule herein described. More particularly, the present invention provides a nucleic acid molecule comprising a sequence of nucleotides substantially as set forth in SEQ ID NO:3 or having at least 15% similarity to all or part thereof or being capable of hybridising under low stringency conditions to a reverse complement of the nucleotide sequence as set forth in SEQ ID NO:3 provided that the

30

- 7 -

nucleic acid sequence having at least 15% similarity but at least 30% dissimilarity to the nucleotide sequence as set forth in SEQ ID NO:3. The nucleotide sequence set forth in SEQ ID NO:3 is also referred to herein as "SOM175". Preferably, the percentage dissimilarity is about 35%, more preferably about 39% and even more preferably about 40-50% or greater.

For the purposes of defining the level of stringency, reference can conveniently be made to Sambrook *et al* (1989) at pages 9.47-9.51 which is herein incorporated by reference where the washing steps disclosed are considered high stringency. A low stringency is defined herein as being in 4-6X SSC/0.1-0.5% w/v SDS at 37-45°C for 2-3 hours. Depending on the source and concentration of nucleic acid involved in the hybridisation, alternative conditions of stringency may be employed such as medium stringent conditions which are considered herein to be 1-4X SSC/0.25-0.5% w/v SDS at $\geq 45^\circ\text{C}$ for 2-3 hours or high stringent conditions considered herein to be 0.1-1X SSC/0.1% w/v SDS at 60°C for 1-3 hours.

The present invention further contemplates a nucleic acid molecule which encodes a VEGF-like molecule as hereinbefore described having at least 15% nucleotide sequence homology to SEQ ID NO:3. Preferred levels of homology include at least about 40%, more preferably around 60-70%.

The present invention is further directed to the murine homologue of human VEGF (referred to herein as "mVRF"). The mVRF has approximately 85% identity and 92% conservation of amino acid residues over the entire coding region compared to human VEGF. The mVRF is encoded by a nucleic acid molecule comprising a nucleotide sequence substantially as set forth in Figure 9.

The VEGF-like molecule of the present invention will be useful in the development of a range of therapeutic and/or diagnostic applications alone or in combination with other molecules such as VEGF. The present invention extends, therefore, to pharmaceutical compositions comprising the VEGF-like molecule or parts, fragments, derivatives, homologues or analogues thereof together with one or more pharmaceutically acceptable

- 8 -

carriers and/or diluents. Furthermore, the present invention extends to vectors comprising the nucleic acid sequence set forth in SEQ ID NO:3 or having at least about 15%, more preferably about 40% and even more preferably around 60-70% similarity thereto but at least 30% and more preferably around 39% dissimilarity thereto and host
5 cells comprising same. In addition, the present invention extends to ribozymes and antisense molecules based on SEQ ID NO:3 as well as neutralizing antibodies to the VEGF-like molecule. Such molecules may be useful in ameliorating the effects of, for example, over expression of VEGF-like genes leading to angiogenesis or vascularization of tumours.

10

Another aspect of the present invention contemplates a method of inducing astroglial proliferation in a mammal, said method comprising administering to said mammal an effective amount of a recombinant proteinaceous molecule having the characteristics:

- 15 (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one property in common with vascular endothelial growth factor (VEGF),

said administration being for a time and under conditions sufficient to induce astroglial
20 proliferation.

Preferably, the recombinant proteinaceous molecule comprises the amino acid sequence set forth in SEQ ID NO:3 or SEQ ID NO:6.

25 A further aspect of the present invention provides a method of promoting neural survival and/or proliferation in a mammal, said method comprising administering to said mammal an effective amount of a recombinant proteinaceous molecule having the characteristics:

- 30 (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one property in common with vascular endothelial growth factor (VEGF),

- 9 -

said administration being for a time and under conditions sufficient to induce astroglial proliferation.

Preferably, the recombinant proteinaceous molecule comprises the amino acid sequence
5 set forth in SEQ ID NO:3 or SEQ ID NO:6.

The present invention also contemplates antibodies to the VEGF-like molecule or nucleic acid probes to a gene encoding the VEGF-like molecule which are useful as diagnostic agents.

10

The present invention is further described by reference to the following non-limiting Figures and/or Examples.

In the Figures:

15

Figure 1 Nucleotide sequence [SEQ ID NO:1] and corresponding amino acid sequence [SEQ ID NO:2] of VEGF₁₆₅.

Figure 2 Nucleotide sequence [SEQ ID NO:3] and corresponding amino acid
20 sequence [SEQ ID NO:4] of SOM175.

Figure 3 Results of BLAST search with SOM175 protein sequence.

Figure 4 BESTFIT alignment of VEGF cDNA and SOM175 cDNA.
25

Figure 5 Multiple alignment of VEGF₁₆₅ with SOM175 and its splice variants at the nucleotide level.

Figure 6 Multiple alignment of VEGF₁₆₅ with SOM175 and its splice variants at
30 the amino acid level.

Figure 7 Diagrammatic representation of SOM175 and its splice variants.

Figure 8(a) Diagrammatic representation of genomic structure of human SOM175 genomic showing exon/intron map.

Figure 8(b) Diagrammatic representation of genomic structure of human SOM175 showing exon/intron boundaries.

Figure 9 Nucleotide and predicted peptide sequences derived from mVRF cDNA clones. Numbering of nucleotides are given on the left, starting from the A of the initiation codon. Amino acids are numbered on the right, starting from the first residue of the predicted mature protein after the putative signal peptide has been removed. The alternately spliced region is double underlined and the resulting peptide sequence from each mRNA is included. A potential polyadenylation signal is indicated in boldface. Start and stop codons of mVRF₁₆₇ and mVRF₁₈₆ are underlined and a polymorphic AC repeat in the 3' UTR is indicated by a stippled box. The positions of intron/exons boundaries are indicated by arrowheads.

Figure 10 BESTFIT alignments of human and murine VRF protein isoforms. A: mVRF₁₆₇ and hVRF₁₆₇. B: mVRF₁₈₆ and hVRF₁₈₆ from the point where the sequences diverge from the respective 167 amino acid isoforms. Amino acid identities are marked with vertical bars and conserved amino acids with colons. An arrow marks the predicted signal peptide cleavage site of human and mouse VRF.

Figure 11 BESTFIT alignment of mVRF₁₆₇ and mVEGF₁₈₈ (Breier *et al*, 1992) peptide sequences. An arrow marks the signal peptide cleavage site of mVEGF. Identical amino acids are indicated by vertical bars and conservative substitutions by colons. Numbering of amino acids is as described in the legend to Figure 9.

Figure 12 Comparison of gene structure between VRF (a generic VRF gene is shown since the intron/exon organisation of the mouse and human homologues is almost identical) and other members of the human VEGF/PIGF/PDGF gene family. Exons are represented by boxes. Protein coding regions and untranslated regions are shown by filled and open sections respectively. The hatched region in VRF indicates the

additional 3' UTR sequence formed by alternate splicing of the VRF₁₈₆ isoform. Potential alternate splice products of each gene are shown.

Figure 13 Autoradiogram of a Northern blot of total RNA from various adult mouse tissues (as indicated) hybridised with an mVRF cDNA clone. A major transcript of 1.3 kb was detected in all samples.

Figure 14 Film autoradiographs (A-C) and dark-field micrographs (D-E) illustrating the expression pattern of mVRF and mRNA in the mouse. In the E14 mouse embryo (A) positive signals are present over the developing heart (Ha) and cerebral cortex (Cx). A low background signal is also present over other tissues in the section. In the E17 embryo (B) and the heart (Ha) is clearly visible due to a strong hybridisation signal. An equally strong signal is present over brown adipose tissue (Fa) in the back and around the thoracic cage. A moderate hybridisation signal is present over the spinal cord (SC) and the tongue (T). The background signal is reduced compared with the E14 embryo. In the young adult mouse (C-D), positive signals are present over the heart (Ha) and adipose tissue (Fa) around the thoracic cage, while, for example, the lungs (Lu) are unlabeled). The hybridisation signal over the heart is evenly distributed over the entire left ventricle, including papillary muscles (D). In the E17 heart hybridised with an excess of cold probe, no positive signal is present (E). Scale bars = 0.5 mm (A), 1.2 mm (B), 1 mm (C), 0.3 mm (D), 0.1 mm (E).

Figure 15 Dark - (A and C) and bright-field (B and D) micrographs showing mVRF mRNA expression in mouse adipose tissue (A-B) and spinal cord (C-D). A strong hybridisation signal is present over fat (A), as shown by the strong labeling in Sudan black stained sections (B). A weak signal is present also in skeletal muscle (M in A-B). In the adult spinal cord (C) the mVRF probes gave a neuronal staining pattern over the gray matter. Toluidine counterstaining showing that motoneurons in the ventral horn (D), interneurons in the deep part of the dorsal horn and around the central canal (not shown) where largely positive for mVRF mRNA. Scale bars = 0.1 mm (A), 0.1 mm (B), 0.25 mm (C), 0.015 mm (D).

- 12 -

Figure 16 Effect of VEGF on embryonic day 8 (E8) chick sensory neurons as determined by % survival, % neurite outgrowth and average neurite length (μm).

Figure 17 Effects of VEGF and SOM175 on chick glia. Tested were CNS glial,
5 peripheral glia and CNS oligodendrocytes.

Figure 18 Effect of various SOM175 proteins on mouse astroglial cells. ■ ^3H (cpm)

1. FGF-2 (10 ng/ml) positive control
2. SOM Δ X6* 1 ng/ml
- 10 3. SOM Δ X6 10 ng/ml
4. SOM Δ X6 100 ng/ml
5. SOM Δ X6 1000 ng/ml
6. SOM Δ X6 1000 ng/ml, no heparin
7. SOMX6** 1 ng/ml
- 15 8. SOMX6 10 ng/ml
9. SOMX6 100 ng/ml
10. SOMX6 1000 ng/ml
11. SOMX6 1000 ng/ml, no heparin

* This refers to SOM175 absent exon 6;

20 ** This refers to SOM175.

Figure 19 Effect of various SOM175 proteins on mouse oligodendroglial cells. ■ ^3H (cpm)

1. FGF-2 (10 ng/ml) positive control
- 25 2. SOM Δ X6* 1 ng/ml
3. SOM Δ X6 10 ng/ml
4. SOM Δ X6 100 ng/ml
5. SOM Δ X6 1000 ng/ml
6. SOM Δ X6 1000 ng/ml, no heparin
- 30 7. SOMX6** 1 ng/ml
8. SOMX6 10 ng/ml
9. SOMX6 100 ng/ml

- 13 -

10. SOMX6 1000 ng/ml

11. SOMX6 1000 ng/ml, no heparin

* This refers to SOM175 absent exon 6;

** This refers to SOM175.

5

Figure 20 Effect of various SOM175 proteins on mouse forebrain neurons. ■ % survival

1. FGF-2 (10 ng/ml) positive control
2. SOM Δ X6* 1 ng/ml
- 10 3. SOM Δ X6 10 ng/ml
4. SOM Δ X6 100 ng/ml
5. SOM Δ X6 1000 ng/ml
6. SOM Δ X6 1000 ng/ml, no heparin
7. SOMX6** 1 ng/ml
- 15 8. SOMX6 10 ng/ml
9. SOMX6 100 ng/ml
10. SOMX6 1000 ng/ml
11. SOMX6 1000 ng/ml, no heparin

* This refers to SOM175 absent exon 6;

20 ** This refers to SOM175.

TABLE 1
SUMMARY OF SEQUENCE IDENTITY NUMBERS

5	SEQ ID NO:1	Nucleotide sequence of VEGF ₁₆₅
	SEQ ID NO:2	Amino acid sequence of VEGF ₁₆₅
	SEQ ID NO:3	Nucleotide sequence of SOM175 (VEGF-like molecules)
	SEQ ID NO:4	Amino acid sequence of SOM175
10	SEQ ID NO:5	Nucleotide sequence of SOM175 absent exon 6
	SEQ ID NO:6	Amino acid sequence of SOM175 absent exon 6
	SEQ ID NO:7	Nucleotide sequence of SOM175 absent exon 6 and exon 7
	SEQ ID NO:8	Amino acid sequence of SOM175 absent exon 6 and exon 7
	SEQ ID NO:9	Nucleotide sequence of SOM175 absent exon 4
15	SEQ ID NO:10	Amino acid sequence of SOM175 absent exon 4
	SEQ ID NO:11	Oligonucleotide
	SEQ ID NO:12	Oligonucleotide
	SEQ ID NO:13	Oligonucleotide
	SEQ ID NO:14	Oligonucleotide

20

EXAMPLE 1

Human cDNA clones

The original SOM175 cDNA was isolated by screening a human foetal brain library
 25 (λzapII, Stratagene) with the cosmid D11S750 (Larsson *et al*, 1992). The plasmid was
 excised "*in vivo*" and a single 1.1kb cDNA was obtained. Three independent SOM175
 cDNAs clones were also isolated from a human foetal spleen library (Stratagene, Uni-
 zap) using the above-mentioned SOM175 insert as a probe. Three clones were obtained:
 SOM175-4A, -5A and -6A. SOM175-5A is an alternately spliced clone with exon 4
 30 being absent (SOM175-e4). These library screens were performed using hybridisation
 conditions recommended by the manufacturer of the library (Stratagene) and random
 primed insert of SOM175.

- 15 -

Two partial human SOM175 cDNAs have also isolated from a λ GT11 human melanoma cell line A2058 library (Clontech) cDNA library screens were performed using hybridisation conditions described by Church and Gilbert, 1984). In each case, the probe was generated by random priming of a PCR product derived from SOM175 (18f-
5 700r).

Mouse cDNA Clones

Human SOM175 was also used to screen a mouse neonatal whole brain cDNA library (Unizap, Stratagene). Four non-chimeric clones were isolated: M175-A, B, C, D. All
10 clones were partial cDNAs and M175-C contained several introns. Three of these cDNAs lacked the exon 6.

Another clone referred to as M1 was completely sequenced and was found to contain the full open reading frame plus part of the 5'utr and total 3'utr.
15

EXAMPLE 2

DNA SEQUENCE ANALYSIS

The entire sequence of the cDNA clone (SOM175) was compiled and is shown in Figure 2 with its corresponding amino acid sequence. This sequence was screened for
20 open reading frames using the MAP program (GCG, University of Wisconsin). A single open reading frame of 672bp was observed (see Figure 2). There appears to be little 5' untranslated sequences (2bp). The 3' untranslated region appears to be complete as it includes a poly-adenylation signal and poly-A tail.

25 Database homology searches were performed using the BLAST algorithm (run at NCBI, USA). This analysis revealed homology to several mammalian forms of VEGF (see Figure 3). The amount of homology between SOM175 and human VEGF₁₆₅ was determined using the BESTFIT program (GCG, University of Wisconsin; see Figures 4 and 5). Nucleotide homology was estimated at 69.7% and protein homology was
30 estimated as at least 33.3% identity and 52.5% conservation using BESTFIT analysis. BLAST analysis on nucleotide sequences revealed the almost complete match to a human expressed sequence tag EST06302 (Adams *et al.*, 1993).

- 16 -

These data indicate that SOM175 encodes a growth factor that has structural similarities to VEGF. Both genes show start and stop codons in similar positions and share discrete blocks of homology. All 8 cysteines as well as a number of other VEGF residues believed to be involved in dimerisation are conserved. These residues are Cysteine-47, 5 Proline-70, Cysteine-72, Valine-74, Arginine-77, Cysteine-78, Glycine-80, Cysteine-81, Cysteine-82, Cysteine-89, Proline-91, Cysteine-122 and Cysteine-124 and are shown in Figure 6. Given the structural conservation between VEGF and the SOM175 gene product it is also possible that they share functional similarities. It is proposed that SOM175 encodes a VEGF-like molecule that shares some properties with VEGF but has 10 unique properties of its own. The nucleotide sequence and corresponding amino acid sequence of VEGF₁₆₅ is shown in Figure 1.

EXAMPLE 3

The percentage similarity and divergence between VEGF₁₆₅ family and SOM175 family 15 (protein) were analysed using the Clustal method, MegAlign Software, DNASTAR, Wisconsin. The results are shown in Tables 2.1 and 2.2. The alternatively spliced forms of SOM175 are abbreviated to SOM715-e6 where all of exon 6 is deleted; SOM715-e6 and 7 where all of exons 6 and 7 are deleted; and SOM175-e4 where all of exon 4 is deleted. The spliced form of SOM175 are shown in Figure 7. Genomic 20 maps of SOM175 showing intron/exon boundaries are shown in Figure 8a and 8b.

Table 2.1

A Percent nucleotide similarity between splice variants of SOM175 and
 5 human VEGF₁₆₅

	VEGF ₁₆₅	SOM175	SOM175-e6	SOM175-e6&7	SOM175-e4
VEGF ₁₆₅	***	34.9	39.7	41.4	37.0
10 SOM175		***	98.9	95.1	99.2
SOM175-e6			***	98.8	84.0
SOM175-e6&7				***	80.3
SOM175-e4					***

15

- 18 -

B Percent nucleotide divergence between splice variants of SOM175 and human VEGF₁₆₅

5		VEGF ₁₆₅	SOM175	SOM175-e6	SOM175-e6&7	SOM175-e4
	VEGF ₁₆₅	***	41.7	41.6	41.7	41.8
	SOM175		***	0.2	0.2	0.0
	SOM175-e6			***	0.0	0.2
10	SOM175-e6&7				***	0.3
	SOM175-e4					***

Table 2.2

15 A Percent amino acid identity between splice variants of SOM175 and human VEGF₁₆₅

		VEGF ₁₆₅	SOM175	SOM175-e6	SOM175-e6&7	SOM175-e4
20	VEGF ₁₆₅	***	31.4	42.3	33.5	40.6
	SOM175		***	74.7	73.7	99.1
	SOM175-e6			***	76.8	99.1
	SOM175-e6&7				***	99.1
	SOM175-e4					***

25

- 19 -

B Percent amino acid divergence between splice variants of SOM175 and human VEGF₁₆₅

5		VEGF ₁₆₅	SOM175	SOM175-e6	SOM175-e6&7	SOM175-e4
	VEGF ₁₆₅	***	65.7	55.4	54.6	57.4
	SOM175		***	19.9	4.2	0.0
	SOM175-e6			***	0.0	0.0
10	SOM175-e6&7				***	0.0
	SOM175-e4					***

15

EXAMPLE 4

BIOASSAYS TO DETERMINE THE FUNCTION OF SOM175

Assays are conducted to evaluate whether SOM175 has similar activities to VEGF on endothelial cell function, angiogenesis and wound healing. Other assays are performed based on the results of receptor binding distribution studies.

20

Assays of endothelial cell function

Endothelial cell proliferation. Endothelial cell growth assays as described in Ferrara & Henzel (1989) and in Gospodarowicz *et al* (1989).

25 *Vascular permeability assay.* This assay, which utilises the Miles test in guinea pigs, will be performed as described in Miles & Miles (1952).

Cell adhesion assay. The influence of SOM175 on adhesion of polymorphs to endothelial cells is analysed.

30

Chemotaxis. This is performed using the standard Boyden chamber chemotaxis assay.

- 20 -

Plasminogen activator assay. Endothelial cells are tested for plasminogen activator and plasminogen activator inhibitor production upon addition of SOM175 (Pepper *et al* (1991)).

- 5 *Endothelial cell migration assay.* The ability of SOM175 to stimulate endothelial cells to migrate and form tubes is assayed as described in Montesano *et al* (1986).

Angiogenesis Assay

- SOM175 induction of an angiogenic response in chick chorioallantoic membrane is
10 evaluated as described in Leung *et al* (1989).

Possible neurotrophic actions of SOM175 are assessed using the following assays:

Neurite outgrowth assay and gene induction (PC12 cells)

- 15 PC12 cells (a phaeochromocytoma cell line) respond to NGF and other neurotrophic factors by developing the characteristics of sympathetic neurons, including the induction of early and late genes and the extension of neurites. These cells are exposed to SOM175 and their response monitored (Drinkwater *et al* (1991); and Drinkwater *et al* (1993)).

20

Cultured neurons from the Peripheral Nervous System (PNS)

Primary cultures of the following PNS neurons are exposed to SOM175 and monitored for any response:

- sensory neurons from neural crest and dorsal root ganglia
- 25 - sympathetic neurons from sympathetic chain ganglia
- placode derived sensory neurons from nodose ganglia
- motoneurons from spinal cord

The assays are described in Suter *et al* (1992) and in Marinou *et al* (1992).

- 30 Where an *in vitro* response is observed, *in vivo* assays for properties such as uptake and retrograde transport are performed as described in Hendry *et al* (1992).

Nerve regeneration (PNS)

Where neurotrophic effects of SOM175 are observed, its possible role in the regeneration of axotomised sensory neurons, sympathetic neurons and motoneurons is analysed by the methods of Otto *et al* (1989); Yip *et al* (1984) and Hendry *et al* (1976).

Actions of SOM175 on CNS neurons

The ability of SOM175 to promote survival of central nervous system neurons is analysed as described in Hagg *et al* (1992); Williams *et al* (1986); Hefti (1986) and Kromer (1987).

Wound Healing

The ability of SOM175 to support wound healing are tested in the most clinically relevant model available, as described in Schilling *et al* (1959) and utilised by Hunt *et al* (1967).

The Haemopoietic System

A variety of *in vitro* and *in vivo* assays on specific cell populations of the haemopoietic system are available and are outlined below:

20 Stem Cells**Murine**

A variety of novel *in vitro* murine stem cell assays have been developed using FACS-purified cells:

25 (a) Repopulating Stem Cells

These are cells capable of repopulating the bone marrow of lethally irradiated mice, and have the Lin⁻, Rh^{hi}, Ly-6A/E⁺, c-kit⁺ phenotype. The test substance is tested on these cells either alone, or by co-incubation with multiple factors, followed by measurement of cellular proliferation by ³H thymidine incorporation.

(b) Late Stage Stem Cells

These are cells that have comparatively little bone marrow repopulating ability but can generate D13 CFU-S. These cells have the Lin⁻, Rh^{hi}, Ly-6A/E⁺, c-kit⁺ phenotype. The test substance is incubated with these cells for a period of time, injected into lethally irradiated recipients, and the number of D13 spleen colonies enumerated.

(c) Progenitor-Enriched Cells

These are cells that respond *in vitro* to single growth factors, and have the Lin⁻, Rh^{hi}, Ly-6A/E⁺, c-kit⁺ phenotype. This assay will show if SOM175 can act directly on haemopoietic progenitor cells. The test substance is incubated with these cells in agar cultures, and the number of colonies enumerated after 7-14 days.

15 Atherosclerosis

Smooth muscle cells play a crucial role in the development or initiation of atherosclerosis, requiring a change in their phenotype from a contractile to a synthetic state. Macrophages, endothelial cells, T lymphocytes and platelets all play a role in the development of atherosclerotic plaques by influencing the growth and phenotypic modulations of smooth muscle cell. An *in vitro* assay that measures the proliferative rate and phenotypic modulations of smooth muscle cells in a multicellular environment is used to assess the effect of SOM175 on smooth muscle cells. The system uses a modified Rose chamber in which different cell types are seeded onto opposite coverslips.

25

Effects of SOM175 on bone

The ability of SOM175 to regulate proliferation of osteoblasts is assayed as described in Lowe *et al* (1991). Any effects on bone resorption are assayed as described in Lowe *et al* (1991). Effects on osteoblast migration and changes in intracellular molecules (e.g. cAMP accumulation, alkaline phosphatase levels) are analysed as described in Midy *et al* (1994).

30

Effects on skeletal muscle cells

Effects of SOM175 on proliferation of myoblasts and development of myotubes can be determined as described by Ewton *et al* (1980) and by Gospodarowicz *et al* (1976).

5

EXAMPLE 5**CLONING MURINE VEGF DNA****Isolation of cDNAs**

Murine VRF (mVRF) clones were selected from a lambda Zap new born whole brain cDNA library (Stratagene). Primary phage from high density filters (5×10^4 pfu/plate) were identified by hybridisation with a 682bp ^{32}P -labelled probe generated by PCR from an hVRF cDNA (pSOM175) as described above. Hybridisation and stringent washes of nylon membranes (Hybond-N) were carried out at 65°C under conditions described by Church and Gilbert (1984). Positive plaques were picked, purified and excised *in vivo* to produce bacterial colonies containing cDNA clones in pBluescript SK-.

Isolation of genomic clones

Genomic clones were isolated from a mouse strain SV/129 library cloned in the lambda Fix II vector (Stratagene). High density filters (5×10^4 pfu/filter) were screened with a 563 bp ^{32}P -labelled probe generated by PCR amplification of the nucleotide 233-798 region of the mVRF cDNA (see Figure 9). Positive clones were plugged and re-screened with filters containing 400-800 pfu. Large scale phage preparations were prepared using the QIAGEN lambda kit or by ZnCl_2 purification (Santos, 1991).

Nucleotide sequencing and analysis

cDNAs were sequenced on both strands using a variety of vector-based and internal primers with Applied Biosystems Incorporated (ABI) dye terminator sequencing kits according to the manufacturer's specifications. Sequences were analysed on an ABI Model 373A automated DNA sequencer. Peptide homology alignments were performed using the program BESTFIT (GCG, Wisconsin).

- 24 -

Identification of intron/exon boundaries

- Identification of exon boundaries and flanking regions was carried out using PCR with mouse genomic DNA or mVRF genomic lambda clones as templates. The primers used in PCR to identify introns were derived from the hVRF sequence and to allow for potential human-mouse sequence mismatches annealing temperatures 5-10°C below the estimated T_m were used. All PCR products were sized by agarose gel electrophoresis and gel purified using QIAquick spin columns (Qiagen) and the intron/exon boundaries were sequenced directly from these products. In addition, some splice junctions were sequenced from subcloned genomic fragments of mVRF.
- 10 Intron/exon boundaries were identified by comparing cDNA and genomic DNA sequences.

Northern analysis

- Total cellular RNA was prepared from a panel of fresh normal adult mouse tissues (brain, kidney, liver, muscle) using the method of Chomczynski and Sacchi (1987). 20µg of total RNA were electrophoresed, transferred to a nylon membrane (Hybond N, Amersham) and hybridised under standard conditions (Church & Gilbert, 1984). Filters were washed at 65°C in 0.1xSSC (20xSSC is 3M NaCl/0.3M trisodium citrate), 0.1% SDS and exposed to X-ray film with intensifying screens at -70°C for 1-3 days.

Characterisation of mVRF cDNAs

- Murine VRF homologues were isolated by screening a murine cDNA library with an hVRF cDNA clone. Five clones of sizes varying from 0.8-1.5 kb were recovered and sequenced. The cDNA sequences were compiled to give a full length 1041 bp cDNA sequence covering the entire open reading frame (621 bp or 564 bp depending on the splice form, see below) and 3' UTR (379 bp), as well as 163 bp of the 5' UTR (Figure 9).
- 30 The predicted initiation codon matched the position of the start codon in hVRF. One other out of frame ATG was located at position -47 and two termination codons were observed upstream (positions -9 and -33, respectively) and in-frame with the putative

- 25 -

initiation codon.

The predicted N-terminal signal peptide of hVRF appears to be present in mVRF with 81% identity (17/21 amino acids). Peptide cleavage within mVRF is expected to occur after residue 21 (Figure 10). These data suggest that mature mVRF is secreted and could therefore conceivably function as a growth factor.

As with hVRF, two open reading frames (ORFs) were detected in cDNAs isolated by library screening. Four of five clones were found to be alternatively spliced and lacked a 101 bp fragment homologous to exon 6 of hVRF. The predicted peptide sequences of the two isoforms of mVRF were determined and aligned with the corresponding human isoforms (Figure 10).

The message encoding mVRF₁₈₆ contains a 621 bp ORF with coding sequences terminating at position +622, towards the end of exon 7 (Figure 9). The smaller message encoding mVRF₁₆₇ actually terminates downstream of the +622 TAG site due to a frame shift resulting from splicing out of the 101 bp exon 6 and the introduction of a stop codon (TGA) at position +666, near the beginning of exon 8 (Figure 9).

The mVRF₁₈₆ protein has strong homology to the amino and central portions of VEGF while the carboxyl end is completely divergent and is alanine rich. mVRF₁₆₇ possesses these similarities and also maintains homology to mVEGF right through to the C-terminus (Figure 11). The overall homology of mVRF₁₆₇ to hVRF₁₆₇ was 85% identity and 92% similarity, respectively (Figure 10). Likewise, homology between mVRF₁₆₇ and mVEGF (Breier *et al*, 1992) was 49% identity and 71% conservative amino acid substitution, respectively (Figure 11).

A canonical vertebrate polyadenylation signal (AATAAA) (Birnstiel *et al*, 1986) was not present in the mVRF cDNA, however, the closely matching sequence GATAAA is present at similar positions in both mouse and human VRF cDNAs (Figure 9). In contrast to hVRF, mVRF was found to contain an AC dinucleotide repeat at the

extreme 3' end of the 3' UTR (nucleotide positions 998 to 1011, Figure 9).

Polymorphism of this repeat region was observed between some of the mVRF cDNAs, with the number of dinucleotides varying from 7 to 11.

5 Genomic characterisation of mVRF

Intron/exon boundaries (Table 3) were mapped using primers which flanked sequences homologous to the corresponding hVRF boundaries. Introns I, III, IV and VI of mVRF (Table 3, Figure 12) were smaller than the hVRF intervening sequences. The complete genomic sequence was compiled from the 5' UTR of mVRF through to intron VI, the largest intervening region (2.2 kb), by sequencing amplified introns and cloned genomic portions of mVRF. There was only one major difference in genomic structure between mVRF and hVRF and that was the exon 7/intron VI boundary of mVRF was located 10bp further downstream in relation to the cDNA sequence, hence exon 7 in mVRF is 10bp longer than the corresponding exon in hVRF.

Exons 6 and 7 are contiguous in mVRF, as has been found to occur in the human homologue. The strong sequence homology between exon 6 of mVRF and hVRF (Figure 10) suggests that this sequence is not a retained intronic sequence but rather encodes a functional part of the VRF₁₈₆ isoform.

General intron/exon structure is conserved between the various members of the VEGF gene family (VEGF, PIGF, hVRF) and therefore it is not surprising that the overall genomic organisation of the mVRF gene is very similar to these genes (Figure 12).

Previous comparative mapping studies have shown that the region surrounding the human multiple endocrine neoplasia type 1 disease locus on chromosome 11q13 is syntenic with the proximal segment of mouse chromosome 19 (Rochelle *et al*, 1992). Since the inventors have mapped the hVRF gene to within 1kb of the human *MEN1* locus (see above) it is most likely that the murine VRF gene maps near the centromere of chromosome 19.

- 27 -

Expression studies of mVRF

Northern analysis of RNA from adult mouse tissues (muscle, heart, lung and liver) showed that expression appears to be ubiquitous and occurs primarily as a major band of approximately 1.3kb in size (Figure 14). This is somewhat different to the pattern observed for hVRF in which two major bands of 2.0 and 5.5 kb have been identified in all tissues examined. The 1.3 kb murine message presumably corresponds to the shorter of the human transcripts and the size variation thereof is most likely due to a difference in the length of the respective 5' UTRs.

10

EXAMPLE 6**EXPRESSION OF MURINE VEGF IN PRE- AND POST-NATAL MOUSE****Animals**

Timed pregnant (n=4) and young adult (n=2) mice (C57 inbred strain, ALAB, Sweden) were sacrificed with carbon dioxide, and the relevant tissues were taken out and frozen on a chuck. Tissues were kept at -70°C until further use. Two gestational ages was used in this study; embryonic day 8 (E8), 14 and E17.

***In situ* hybridisation histochemistry**

In situ hybridisation was performed as previously described (Dagerlind *et al*, 1992).

Briefly, transverse sections (14µm) were cut in a cryostat (Microm, Germany), thawed onto Probe-On slides (Fisher Scientific, USA) and stored in black sealed boxes at -70°C until used. The sequences of the synthetic 42-mer oligonucleotides complementary to mRNA encoding mVRF were

ACCACCACCTCCCTGGGCTGGCATGTGGCACGTGCATAAACG [SEQ ID NO:11] (complementary to nt 120-161) and

AGTTGTTTGACCACATTGCCCATGAGTTCCATGCTCAGAGGC [SEQ ID NO:12] (complementary to nt 162-203). To detect the two alternative splice forms oligonucleotide GATCCTGGGGCTGGAGTGGGATGGATGATGTCAGCTGG [SEQ ID NO:13] (complementary to nt xxx-xxx) and

GCGGGCAGAGGATCCTGGGGCTGTCTGGCCTCACAGCACT [SEQ ID NO:14] were used. The probes were labeled at the 3'-end with deoxyadenosine-alpha[thio]triphosphate [³⁵S] (NEN, USA) using terminal deoxynucleotidyl

- 28 -

transferase (IBI, USA) to a specific activity of $7-10 \times 10^8$ cpm/ μ g and hybridised to the sections without pretreatment for 16-18 h at 42°C. The hybridisation mixture contained: 50% v/v formamide, 4 x SSC (1 x SSC = 0.15M NaCl and 0.015M sodium-citrate), 1 x Denhardt's solution (0.02% each of polyvinyl-pyrrolidone, BSA and Ficoll), 1% v/v sarcosyl (N-lauroylsarcosine; Sigma), 0.02M phosphate buffer (pH 7.0), 10% w/v dextran sulfate (Pharmacia, Sweden), 250 μ g/ml yeast tRNA (Sigma), 500 μ g/ml sheared and heat denatured salmon sperm DNA (Sigma) and 200mM dithiothreitol (DTT; LKB, Sweden). In control sections, the specificity of both probes was checked by adding a 20-fold excess of unlabeled probe to the hybridisation mixture. In addition, adjacent sections were hybridised with a probe unrelated to this study which gave a different expression pattern. Following hybridisation the sections were washed several times in 1 x SSC at 55°C, dehydrated in ethanol and dipped in NTB2 nuclear track emulsion (Kodak, USA). After 3-5 weeks the sections were development in D-19 developer (Kodak, USA) and cover-slipped. In some cases, sections were opposed to an autoradiographic film (Beta-max autoradiography film Amersham Ltd, UK) prior to emulsion-dipping.

The four different probes gave identical hybridisation patterns in all tissues examined. Mouse VRF expression was detecting already in the E8 embryo, in which positive signal was recorded over structures most likely corresponding to the neuronal tube. In sagittal sections of E14 mouse embryo the strongest hybridisation signal was present over heart and in the nervous system, especially cerebral cortex (Figure 14A). A low level of expression was present in all other tissues. At a later gestational age, E17, a high mVRF mRNA signal was confined to the heart and brown fat tissue in the back and around the neck (Figure 14B). Clearly positive hybridisation signals were present in the gray of the spinal cord and in the tongue (Figure 14B). Expression in the cerebral cortex was clearly reduced compared to day 14. The weak background expression seen in the E14 embryo in for example muscle, had decreased at this gestational age. A strong mVRF mRNA hybridisation signal was present solely over the heart and in the brown fat in the young adult mice (Figure 14C). The signal over the heart was evenly distributed over the entire ventricular wall, including the papillary muscles (Figure 14D). In sections of heart tissue hybridised with an

- 29 -

excess of cold probe, no specific labeling over background signal was recorded (Figure 14E).

Apart from the heart, mVRF mRNA signal was present over certain tissues on the
5 outside of the thoracic cage that morphologically resembled brown fat. This was
verified with sudan black counterstaining, which showed a strong staining in the
same areas (Figure 15A and 15B). In transverse sections of adult mouse spinal cord,
the mVRF probes gave a neuronal staining pattern over the gray matter (Figure 15C).
Counterstaining with toluidine (Figure 15D) showed that motoneurons in the ventral
10 horn (Figure 15C and 15D), interneurons (Figure 15C) in the deep part of the dorsal
horn and around the central canal where to a large extent positive for mVRF mRNA.

EXAMPLE 7

EFFECTS OF VEGF AND SOM175 PROTEINS ON CHICK

15 SENSORY NEURONS

The effects of VEGF and SOM175 proteins on embryonic day 8 chick sensory
neurons were determined using the method of Nurcombe *et al* (1992). The neuronal
assay was read at 48 hours using 2000 cells per assay well. The results were
obtained using ³H-thymidine counts. The percentage survival of neurons, neurite
20 outgrowth and average neurite length in μm were determined using NGF as positive
control and various concentrations of VEGF, VEGF in the presence of heparin and
VEGF in the presence of heparin and 5 μM , 5'-flurouracil (5FU). 5FU kills glial
cells.

25 The results are shown in Figure 16. The results show that VEGF is effective in
promoting neuronal survival but that this requires the presence of glial cells. Figure
17 shows the results of the effect of VEGF and SOM175 on three types of chick
glia. The glia tested were CNS glia, peripheral glia and CNS oligodendrocytes.
Heparin was used as 10 $\mu\text{g/ml}$ in all cultures and the assay was read at 24 hours.
30 Results were measured in ³H-thymidine counts using 2000 cells per well.

- 30 -

The results show that for chick central and peripheral neurons, astroglia were markedly stimulated to proliferate by SOM175 in the presence of heparin but that chick oligodendrocytes showed negligible increase in the rate of division.

5

EXAMPLE 8

EFFECTS OF SOM175 PROTEINS ON MOUSE PRIMARY AND CENTRAL NEURONS

The results in Example 7 show that VEGF isoform had an effect on chick primary and central neurons through the agency of the astroglial cells. Similar experiments were repeated in mouse cells.

Culture conditions

Neuronal and glial cells for all *in vitro* experiments were prepared and cultured according to the techniques described in "Methods in Neurosciences (Vol. 2): Cell Culture" Ed. P.M. Conn, Academic Press, San Diego, 1990, pp33-46 for astroglial cells, pp56-74 for oligodendroglial cells, and pp87-102 for central neurons.

Cells were plated onto 24-well culture clusters (Nunc) coated with poly-L-ornithine (0.1 mg/ml, 1h) at a density of 2,000 cells/well. After 48 hours in culture, neurons were counted in the wells under inverted phase light using well established techniques (Maruta *et al.* 1993) and glial cells assessed with [³H]thymidine uptake to monitor cell division rates as below. Heparin (10µg/ml, low molecular weight fraction, Sigma Chemical Corp.) was present at all times in the culture media except where noted. The neuronal cultures were supplemented with 5mM 5-fluoro-2-deoxyuridine (Sigma) to suppress background glial growth.

³H-Thymidine incorporation assay for glial cell proliferation

The cells were pulsed for 14h with ³H-thymidine (specific activity 103 µCi/ug) from a stock concentration of 0.1 mCi/ml in standard medium, giving a final incubating volume of 20 µl/well. The contents of the wells were harvested and absorbed onto nitrocellulose paper (Titertek, Flow). Remaining adherent cells were removed by

- 31 -

incubation with trypsin/versene (CSL Limited, Victoria, Australia) for 5 min. This procedure was carried out twice. The nitrocellulose discs were washed in a standard Titertek harvester (Flow) using first distilled water, and then methanol. The nitrocellulose discs were dried, scintillation fluid (containing 5% v/v Triton-X) added and the discs counted on a scintillation counter.

Greatest activity was seen with preparations of SOM175 absent exon 6 (SOM Δ X6) on mouse astroglial cell cultures, where there was a significant stimulus to their proliferation when delivered in conjunction with heparin (Figure 16). Little stimulus was given to the proliferation of oligodendroglial cells (Figure 17), and very little discernable potentiation of the survival response of isolated forebrain neurons (Figure 18). The standard deviation on all three graphs for each point was less than 8%.

The viability of neurons can be maintained by promoting glial cell proliferation. Furthermore, SOM Δ X6 is a good inducer of astroglial proliferation and may be expressed in conjunction with the formation of astroglial endfeet on central nervous system endothelial cells.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

TABLE 3
Splice junctions of the murine VRF gene

5	5' UTR*	Exon 1 >223bp	CCCAGgtacgtgcgt	Intron I	495bp
	ttccccacagGCCCC	Exon 2 43bp	GAAAGgtaataatag	Intron II	288bp
	ctgcccacagTGGTG	Exon 3 197bp	TGCAGgtaccagggc	Intron III	196bp
	ctgagcacagATCCT	Exon 4 74bp	TGCAGgtgccagccc	Intron IV	182bp
	ctcttttcagACCTA	Exon 5 36bp	GACAGattcttggtg	Intron V	191bp
10	ctcctcctagGGTTG	Exon 6 101bp		(no intron)	
	CCCACTCCAGCCCCA	Exon 7 135bp	TGTAGgtaaggagtc	Intron VI	~2200bp
	cactccccagGTGCC	Exon 8 394bp	AGAGATGGAGACACT		

Uppercase and lowercase letters denote exonic and intronic sequences respectively.

15 * Indicates that the 5' end of exon 1 has not yet been determined.

BIBLIOGRAPHY

- Adams MD, Soares MB, Kerlavage AR, Fields C, Venter JC, (1993) *Nature Genet.* 4, 373-380.
- Birnstiel ML, Busslinger M and Strub K (1985) *Cell* 41, 349-359.
- Breier G, Albrecht U, Sterrer S and Risau W (1992) *Development* 114, 521-532.
- Chomczynski P and Sacchi N (1987) *Analyt. Biochem.* 162, 156-159.
- Church G and Gilbert W (1984) *Proc. Natl. Acad. Sci. USA* 81, 1991-1995.
- Dagerlind A, Friberg K, Bean AJ and Hokfelt T (1992) *Histochemistry* 98, 39-49.
- Dissen GA, Lara HE, Fabrenbach WH, Costa ME, Ojeda SR, (1994) *Endocrinology* 134, 1146-1154.
- Drinkwater CC, Barker PA, Suter U and Shooter EM (1993) *J. Biol. Chem.*, 268, 23202-23207.
- Drinkwater CC, Suter U, Angst C and Shooter EM (1991) *Proc. Roy. Soc. Lond. (Series B)*, 246, 307-313.
- Ewton DZ & Florini JR (1980) *Endocrinology*, 106: 577-583.
- Ferrara N & Henzel WJ (1989) *Biochem. Biophys. Res. Commun.* 161, 851-858.
- Folkman J & Shing Y (1992) *J. Biol. Chem.* 267, 10931-10934.
- Gospodarowicz D, Abraham JA & Schilling J (1989) *Proc. Natl. Acad. Sci USA* 86, 7311-7315.
- Gospodarowicz D, Weseman J, Morgan JS & Lindstrom J (1976) *J. Cell Biol.*, 70: 395-405.
- Hagg T, Quon D, Higaki J & Varon S (1992) *Neuron*, 8, 145-158.
- Hefti S (1986) *J. Neurosci.*, 6, 2155-2162.
- Hendry IA & Campbell J (1976) *J. Neurocytol.*, 5, 351-360.
- Hendry IA, Murphy M, Hilton DJ, Nicola NA & Bartlett PF (1992) *J. Neurosci.* 12, 3427-3434.
- Hunt *et al.*, (1967) *Am. J. Surgery*, 114: 302-307.
- Koch AE, Harlow LA, Haines GK, Amento EP, Unemoti EN, Wong WL, Pope RM, Ferrara N, (1994) *J. Immunol.* 152, 4149-4156.
- Kromer AF (1987) *Science*, 235, 214-216.

- Larsson C, Weber G, Kvanta E, Lewis C, Janson M, Jones C, Glaser T, Evans G, Nordenskjold M, (1992) *Hum. Genet.* 89, 187-193.
- OZUS Leung DW, Cachianes G, Kuang W-J, Goeddel DV & Ferrara N (1989) *Science* 246:1306-1309.
- Lowe C, Cornish J, Callon K, Martin TJ & Reid IR (1991) *J. Bone Mineral Res.*, 6, 1277-1283.
- Lowe C, Cornish J, Martin TJ & Reid IR (1991) *Calcif. Tissue Int.*, 49, 394-397.
- Martinou JC, Martinou I & Kato AC (1992) *Neuron*, 8, 737-744.
- Maruta *et al* (1993) *Growth Factors* 8: 119-134.
- Midy V & Plouet J (1994) *Biochem. Biophys. Res. Commun.*, 199: 380-386.
- Miles AA & Miles EM (1952) *J. Physiol. (Lond)* 118:228-257.
- Montesano R, Vassalli JD, Baird A, Guillemin R & Orci, L (1986) *Proc. Natl. Acad. Sci USA*, 83, 7297-7301.
- Nurcombe *et al* (1992) *Development* 116: 1175-1183.
- Otto D., Frotscher M & Unsicker K (1989) *J. Neurosci. Res.*, 22, 83-91.
- Pepper MS, Ferrara N, Orci L, Montesano R. (1991) *Biochem. Biophys. Res. Commun.* 181, 902-906).
- Rochell JM, Watson ML, Oakey RJ and Seldin MF (1992) *Genomics* 14, 26-31.
- Roth S & Weston J (1967) *Proc. Natl. Acad. Sci USA*, 58: 974-980.
- Sambrook J, Fritsch EF, Maniatis T, (1989) *Molecular Cloning: A Laboratory Manual - 2nd Ed.* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Santos MA (1991) *Nucleic Acids Res.* 19, 5442.
- Schilling *et al.*, (1959) *Surgery*, 46: 702-710.
- Senger DR, Van De Water L, Brown LF, Nagy JA, Yeo KT, Yeo TK, Berse B, Jackman RW, Dvorak AM, Dvorak HF (1993) *Cancer Metastasis Rev.* 12, 303-324.
- Sharkey AM, Charnock-Jones DS, Boocock CA, Brown KD, Smith SK, (1993) *J. Reprod. Fertil.* 99, 609-615.
- Sunderkotter C, Steinbrink K, Goebeler M, Bhardway R, Sorg E, (1993) *J. Leukocyt. Biol.* 55, 410-422.
- Suter U, Angst C, Tien C-L, Drinkwater CC, Lindsay RM and Shooter EM (1992) *J. Neurosci.*, 12, 306-318.

Tischer E, Mitchell R, Hartman T, Silva M, Gospodarowicz D, Fiddes JC, & Abraham J (1991) *J. Biol. Chem.* 266, 11947-11954.

Williams LR, Varon S, Peterson GM, Victorin K, Fischer W, Bjorklund A & Gage FH (1986) *Proc. Natl. Acad. Sci. USA* 83, 9231-9235.

Yan Z, Weich HA, Bernart W, Breckwoldt M, Neulen J, (1993) *J. Clin. Endocrinol. Metab.* 77, 1723-1725.

Yip NK, Rich KM, Lampe PA & Johnson EM Jr (1984) *J. Neurosci.*, 4, 2986-2992.

- 36 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(countries other than US) AMRAD OPERATIONS PTY. LTD.
(us only) Hayward, N and Weber, G

(ii) TITLE OF INVENTION: A NOVEL GROWTH FACTOR AND A
GENETIC SEQUENCE ENCODING SAME

(iii) NUMBER OF SEQUENCES: 14

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: DAVIES COLLISON CAVE
(B) STREET: 1 LITTLE COLLINS STREET
(C) CITY: MELBOURNE
(D) STATE: VICTORIA
(E) COUNTRY: AUSTRALIA
(F) ZIP: 3000

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: PCT INTERNATIONAL
(B) FILING DATE: 22-FEB-1996

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: AU PN1457
(B) FILING DATE: 02-MAR-1995
(A) APPLICATION NUMBER: AU PN6647
(B) FILING DATE: 20-NOV-1995
(A) APPLICATION NUMBER: AU PN7274
(B) FILING DATE: 22-DEC-1995

- 37 -

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: HUGHES DR, E JOHN L

(C) REFERENCE/DOCKET NUMBER: EJH/EK

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: +61 3 9254 2777

(B) TELEFAX: +61 3 9254 2770

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 649 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 17...589

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCGGGCCTCC GAAACC ATG AAC TTT CTG CTG TCT TGG GTG CAT TGG AGC	49
Met Asn Phe Leu Leu Ser Trp Val His Trp Ser	
1 5 10	
CTT GCC TTG CTG CTC TAC CTC CAC CAT GCC AAG TGG TCC CAG GCT GCA	97
Leu Ala Leu Leu Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala	
15 20 25	
CCC ATG GCA GAA GGA GGA GGG CAG AAT CAT CAC GAA GTG GTG AAG TTC	145
Pro Met Ala Glu Gly Gly Gly Gln Asn His His Glu Val Val Lys Phe	
30 35 40	
ATG GAT GTC TAT CAG CGC AGC TAC TGC CAT CCA ATC GAG ACC CTG GTG	193
Met Asp Val Tyr Gln Arg Ser Tyr Cys His Pro Ile Glu Thr Leu Val	
45 50 55	
GAC ATC TTC CAG GAG TAC CCT GAT GAG ATC GAG TAC ATC TTC AAG CCA	241
Asp Ile Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro	
60 65 70 75	
TCC TGT GTG CCC CTG ATG CGA TGC GGG GGC TGC TGC AAT GAC GAG GGC	289
Ser Cys Val Pro Leu Met Arg Cys Gly Gly Cys Cys Asn Asp Glu Gly	
80 85 90	

- 38 -

CTG GAG TGT GTG CCC ACT GAG GAG TCC AAC ATC ACC ATG CAG ATT ATG Leu Glu Cys Val Pro Thr Glu Glu Ser Asn Ile Thr Met Gln Ile Met 95 100 105	337
CGG ATC AAA CCT CAC CAA GGC CAG CAC ATA GGA GAG ATG AGC TTC CTA Arg Ile Lys Pro His Gln Gly Gln His Ile Gly Glu Met Ser Phe Leu 110 115 120	385
CAG CAC AAC AAA TGT GAA TGC AGA CCA AAG AAA GAT AGA GCA AGA CAA Gln His Asn Lys Cys Glu Cys Arg Pro Lys Lys Asp Arg Ala Arg Gln 125 130 135	433
GAA AAT CCC TGT GGG CCT TGC TCA GAG CGG AGA AAG CAT TTG TTT GTA Glu Asn Pro Cys Gly Pro Cys Ser Glu Arg Arg Lys His Leu Phe Val 140 145 150 155	481
CAA GAT CCG CAG ACG TGT AAA TGT TCC TGC AAA AAC ACA GAC TCG CGT Gln Asp Pro Gln Thr Cys Lys Cys Ser Cys Lys Asn Thr Asp Ser Arg 160 165 170	529
TGC AAG GCG AGG CAG CTT GAG TTA AAC GAA CGT ACT TGC AGA TGT GAC Cys Lys Ala Arg Gln Leu Glu Leu Asn Glu Arg Thr Cys Arg Cys Asp 175 180 185	577
AAG CCG AGG CGG TGAGCCGGGC AGGAGGAAGG AGCCTCCCTC AGCGTTTCGG Lys Pro Arg Arg 190	629
GAACCAGATC TCTCACCAGG	649

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 191 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Asn Phe Leu Leu Ser Trp Val His Trp Ser Leu Ala Leu Leu Leu 1 5 10 15
Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Met Ala Glu Gly 20 25 30
Gly Gly Gln Asn His His Glu Val Val Lys Phe Met Asp Val Tyr Gln 35 40 45
Arg Ser Tyr Cys His Pro Ile Glu Thr Leu Val Asp Ile Phe Gln Glu 50 55 60

- 39 -

Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser Cys Val Pro Leu
 65 70 75 80
 Met Arg Cys Gly Gly Cys Cys Asn Asp Glu Gly Leu Glu Cys Val Pro
 85 90 95
 Thr Glu Glu Ser Asn Ile Thr Met Gln Ile Met Arg Ile Lys Pro His
 100 105 110
 Gln Gly Gln His Ile Gly Glu Met Ser Phe Leu Gln His Asn Lys Cys
 115 120 125
 Glu Cys Arg Pro Lys Lys Asp Arg Ala Arg Gln Glu Asn Pro Cys Gly
 130 135 140
 Pro Cys Ser Glu Arg Arg Lys His Leu Phe Val Gln Asp Pro Gln Thr
 145 150 155 160
 Cys Lys Cys Ser Cys Lys Asn Thr Asp Ser Arg Cys Lys Ala Arg Gln
 165 170 175
 Leu Glu Leu Asn Glu Arg Thr Cys Arg Cys Asp Lys Pro Arg Arg
 180 185 190

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1094 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 3..624

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CC ATG AGC CCT CTG CTC CGC CGC CTG CTG CTC GCC GCA CTC CTG CAG 47
 Met Ser Pro Leu Leu Arg Arg Leu Leu Leu Ala Ala Leu Leu Gln
 1 5 10 15
 CTG GCC CCC GCC CAG GCC CCT GTC TCC CAG CCT GAT GCC CCT GGC CAC 95
 Leu Ala Pro Ala Gln Ala Pro Val Ser Gln Pro Asp Ala Pro Gly His
 20 25 30
 CAG AGG AAA GTG GTG TCA TGG ATA GAT GTG TAT ACT CGC GCT ACC TGC 143
 Gln Arg Lys Val Val Ser Trp Ile Asp Val Tyr Thr Arg Ala Thr Cys
 35 40 45

- 40 -

CAG CCC CGG GAG GTG GTG GTG CCC TTG ACT GTG GAG CTC ATG GGC ACC Gln Pro Arg Glu Val Val Val Pro Leu Thr Val Glu Leu Met Gly Thr 50 55 60	191
GTG GCC AAA CAG CTG GTG CCC AGC TGC GTG ACT GTG CAG CGC TGT GGT Val Ala Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly 65 70 75	239
GGC TGC TGC CCT GAC GAT GGC CTG GAG TGT GTG CCC ACT GGG CAG CAC Gly Cys Cys Pro Asp Asp Gly Leu Glu Cys Val Pro Thr Gly Gln His 80 85 90 95	287
CAA GTC CGG ATG CAG ATC CTC ATG ATC CGG TAC CCG AGC AGT CAG CTG Gln Val Arg Met Gln Ile Leu Met Ile Arg Tyr Pro Ser Ser Gln Leu 100 105 110	335
GGG GAG ATG TCC CTG GAA GAA CAC AGC CAG TGT GAA TGC AGA CCT AAA Gly Glu Met Ser Leu Glu Glu His Ser Gln Cys Glu Cys Arg Pro Lys 115 120 125	383
AAA AAG GAC AGT GCT GTG AAG CCA GAC AGG GCT GCC ACT CCC CAC CAC Lys Lys Asp Ser Ala Val Lys Pro Asp Arg Ala Ala Thr Pro His His 130 135 140	431
CGT CCC CAG CCC CGT TCT GTT CCG GGC TGG GAC TCT GCC CCC GGA GCA Arg Pro Gln Pro Arg Ser Val Pro Gly Trp Asp Ser Ala Pro Gly Ala 145 150 155	479
CCC TCC CCA GCT GAC ATC ACC CAT CCC ACT CCA GCC CCA GGC CCC TCT Pro Ser Pro Ala Asp Ile Thr His Pro Thr Pro Ala Pro Gly Pro Ser 160 165 170 175	527
GCC CAC GCT GCA CCC AGC ACC ACC AGC GCC CTG ACC CCC GGA CCT GCC Ala His Ala Ala Pro Ser Thr Thr Ser Ala Leu Thr Pro Gly Pro Ala 180 185 190	575
GCT GCC GCT GCC GAC GCC GCA GCT TCC TCC GTT GCC AAG GGC GGG GCT T Ala Ala Ala Ala Asp Ala Ala Ala Ser Ser Val Ala Lys Gly Gly Ala 195 200 205	624
AGAGCTCAAC CCAGACACCT GCAGGTGCCG GAAGCTGCGA AGGTGACACA TGGCTTTTCA	684
GACTCAGCAG GGTGACTTGC CTCAGAGGCT ATATCCCACT GGGGGAACAA AGGGGAGCCT	744
GGTAAAAAAC AGCCAAGCCC CCAAGACCTC AGCCCAGGCA GAAGCTGCTC TAGGACCTGG	804
GCCTCTCAGA GGGCTCTTCT GCCATCCCTT GTCTCCCTGA GGCCATCATC AAACAGGACA	864
GAGTTGGAAG AGGAGACTGG GAGGCAGCAA GAGGGGTCAC ATACCAGCTC AGGGGAGAAT	924
GGAGTACTGT CTCAGTTTCT AACCACCTCTG TGCAAGTAAG CATCTTACAA CTGGCTCTTC	984
CTCCCCTCAC TAAGAAGACC CAAACCTCTG CATAATGGGA TTTGGGCTTT GGTACAAGAA	1044
CTGTGACCCC CAACCCTGAT AAAAGAGATG GAAGGAAAAA AAAAAAAAAA	1094

- 41 -

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 207 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Ser Pro Leu Leu Arg Arg Leu Leu Leu Ala Ala Leu Leu Gln Leu
 1             5             10             15

Ala Pro Ala Gln Ala Pro Val Ser Gln Pro Asp Ala Pro Gly His Gln
      20             25             30

Arg Lys Val Val Ser Trp Ile Asp Val Tyr Thr Arg Ala Thr Cys Gln
      35             40             45

Pro Arg Glu Val Val Val Pro Leu Thr Val Glu Leu Met Gly Thr Val
      50             55             60

Ala Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly Gly
      65             70             75             80

Cys Cys Pro Asp Asp Gly Leu Glu Cys Val Pro Thr Gly Gln His Gln
      85             90             95

Val Arg Met Gln Ile Leu Met Ile Arg Tyr Pro Ser Ser Gln Leu Gly
      100            105            110

Glu Met Ser Leu Glu Glu His Ser Gln Cys Glu Cys Arg Pro Lys Lys
      115            120            125

Lys Asp Ser Ala Val Lys Pro Asp Arg Ala Ala Thr Pro His His Arg
      130            135            140

Pro Gln Pro Arg Ser Val Pro Gly Trp Asp Ser Ala Pro Gly Ala Pro
      145            150            155            160

Ser Pro Ala Asp Ile Thr His Pro Thr Pro Ala Pro Gly Pro Ser Ala
      165            170            175

His Ala Ala Pro Ser Thr Thr Ser Ala Leu Thr Pro Gly Pro Ala Ala
      180            185            190

Ala Ala Ala Asp Ala Ala Ala Ser Ser Val Ala Lys Gly Gly Ala
      195            200            205

```

- 42 -

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 993 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 3..566

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

CC ATG AGC CCT CTG CTC CGC CGC CTG CTG CTC GCC GCA CTC CTG CAG      47
Met Ser Pro Leu Leu Arg Arg Leu Leu Leu Ala Ala Leu Leu Gln
      1              5              10              15

CTG GCC CCC GCC CAG GCC CCT GTC TCC CAG CCT GAT GCC CCT GGC CAC      95
Leu Ala Pro Ala Gln Ala Pro Val Ser Gln Pro Asp Ala Pro Gly His
      20              25              30

CAG AGG AAA GTG GTG TCA TGG ATA GAT GTG TAT ACT CGC GCT ACC TGC      143
Gln Arg Lys Val Val Ser Trp Ile Asp Val Tyr Thr Arg Ala Thr Cys
      35              40              45

CAG CCC CGG GAG GTG GTG GTG CCC TTG ACT GTG GAG CTC ATG GGC ACC      191
Gln Pro Arg Glu Val Val Val Pro Leu Thr Val Glu Leu Met Gly Thr
      50              55              60

GTG GCC AAA CAG CTG GTG CCC AGC TGC GTG ACT GTG CAG CGC TGT GGT      239
Val Ala Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly
      65              70              75

GGC TGC TGC CCT GAC GAT GGC CTG GAG TGT GTG CCC ACT GGG CAG CAC      287
Gly Cys Cys Pro Asp Asp Gly Leu Glu Cys Val Pro Thr Gly Gln His
      80              85              90              95

CAA GTC CGG ATG CAG ATC CTC ATG ATC CGG TAC CCG AGC AGT CAG CTG      335
Gln Val Arg Met Gln Ile Leu Met Ile Arg Tyr Pro Ser Ser Gln Leu
      100              105              110

GGG GAG ATG TCC CTG GAA GAA CAC AGC CAG TGT GAA TGC AGA CCT AAA      383
Gly Glu Met Ser Leu Glu Glu His Ser Gln Cys Glu Cys Arg Pro Lys
      115              120              125

AAA AAG GAC AGT GCT GTG AAG CCA GAT AGC CCC AGG CCC CTC TGC CCA      431
Lys Lys Asp Ser Ala Val Lys Pro Asp Ser Pro Arg Pro Leu Cys Pro
      130              135              140

```

- 43 -

CGC TGC ACC CAG CAC CAC CAG CGC CCT GAC CCC CGG ACC TGC CGC TGC 479
 Arg Cys Thr Gln His His Gln Arg Pro Asp Pro Arg Thr Cys Arg Cys
 145 150 155

CGC TGC CGA CGC CGC AGC TTC CTC CGT TGC CAA GGG CGG GGC TTA GAG 527
 Arg Cys Arg Arg Arg Ser Phe Leu Arg Cys Gln Gly Arg Gly Leu Glu
 160 165 170 175

CTC AAC CCA GAC ACC TGC AGG TGC CGG AAG CTG CGA AGG TGACACATGG 576
 Leu Asn Pro Asp Thr Cys Arg Cys Arg Lys Leu Arg Arg
 180 185

CTTTTCAGAC TCAGCAGGGT GACTTGCCTC AGAGGCTATA TCCCAGTGGG GGAACAAAGG 636

GGAGCCTGGT AAAAAACAGC CAAGCCCCCA AGACCTCAGC CCAGGCAGAA GCTGCTCTAG 696

GACCTGGGCC TCTCAGAGGG CTCTTCTGCC ATCCCTTGTC TCCCTGAGGC CATCATCAAA 756

CAGGACAGAG TTGGAAGAGG AGACTGGGAG GCAGCAAGAG GGGTCACATA CCAGCTCAGG 816

GGAGAATGGA GTACTGTCTC AGTTTCTAAC CACTCTGTGC AAGTAAGCAT CTTACAACCTG 876

GCTCTTCCTC CCCTCACTAA GAAGACCCAA ACCTCTGCAT AATGGGATTT GGGCTTTGGT 936

ACAAGAACTG TGACCCCCAA CCCTGATAAA AGAGATGGAA GGAAAAAAAA AAAAAAA 993

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 188 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ser Pro Leu Leu Arg Arg Leu Leu Leu Ala Ala Leu Leu Gln Leu
 1 5 10 15

Ala Pro Ala Gln Ala Pro Val Ser Gln Pro Asp Ala Pro Gly His Gln
 20 25 30

Arg Lys Val Val Ser Trp Ile Asp Val Tyr Thr Arg Ala Thr Cys Gln
 35 40 45

Pro Arg Glu Val Val Val Pro Leu Thr Val Glu Leu Met Gly Thr Val
 50 55 60

Ala Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly Gly
 65 70 75 80

- 44 -

Cys	Cys	Pro	Asp	Asp	Gly	Leu	Glu	Cys	Val	Pro	Thr	Gly	Gln	His	Gln	85	90	95
Val	Arg	Met	Gln	Ile	Leu	Met	Ile	Arg	Tyr	Pro	Ser	Ser	Gln	Leu	Gly	100	105	110
Glu	Met	Ser	Leu	Glu	Glu	His	Ser	Gln	Cys	Glu	Cys	Arg	Pro	Lys	Lys	115	120	125
Lys	Asp	Ser	Ala	Val	Lys	Pro	Asp	Ser	Pro	Arg	Pro	Leu	Cys	Pro	Arg	130	135	140
Cys	Thr	Gln	His	His	Gln	Arg	Pro	Asp	Pro	Arg	Thr	Cys	Arg	Cys	Arg	145	150	155
Cys	Arg	Arg	Arg	Ser	Phe	Leu	Arg	Cys	Gln	Gly	Arg	Gly	Leu	Glu	Leu	165	170	175
Asn	Pro	Asp	Thr	Cys	Arg	Cys	Arg	Lys	Leu	Arg	Arg					180	185	

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 858 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) **FEATURE:**

- (A) NAME/KEY: CDS
(B) LOCATION: 3..431

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CC	ATG	AGC	CCT	CTG	CTC	CGC	CGC	CTG	CTC	GCC	GCA	CTC	CTG	CAG	47	
Met	Ser	Pro	Leu	Leu	Arg	Arg	Leu	Leu	Leu	Ala	Ala	Leu	Leu	Gln		
1					5					10				15		
CTG	GCC	CCC	GCC	CAG	GCC	CCT	GTC	TCC	CAG	CCT	GAT	GCC	CCT	GGC	CAC	95
Leu	Ala	Pro	Ala	Gln	Ala	Pro	Val	Ser	Gln	Pro	Asp	Ala	Pro	Gly	His	
				20					25					30		
CAG	AGG	AAA	GTG	GTG	TCA	TGG	ATA	GAT	GTG	TAT	ACT	CGC	GCT	ACC	TGC	143
Gln	Arg	Lys	Val	Val	Ser	Trp	Ile	Asp	Val	Tyr	Thr	Arg	Ala	Thr	Cys	
			35					40						45		
CAG	CCC	CGG	GAG	GTG	GTG	GTG	CCC	TTG	ACT	GTG	GAG	CTC	ATG	GGC	ACC	191
Gln	Pro	Arg	Glu	Val	Val	Val	Pro	Leu	Thr	Val	Glu	Leu	Met	Gly	Thr	
			50				55					60				

- 45 -

GTG GCC AAA CAG CTG GTG CCC AGC TGC GTG ACT GTG CAG CGC TGT GGT 239
Val Ala Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly
65 70 75

GGC TGC TGC CCT GAC GAT GGC CTG GAG TGT GTG CCC ACT GGG CAG CAC 287
Gly Cys Cys Pro Asp Asp Gly Leu Glu Cys Val Pro Thr Gly Gln His
80 85 90 95

CAA GTC CGG ATG CAG ATC CTC ATG ATC CGG TAC CCG AGC AGT CAG CTG 335
Gln Val Arg Met Gln Ile Leu Met Ile Arg Tyr Pro Ser Ser Gln Leu
100 105 110

GGG GAG ATG TCC CTG GAA GAA CAC AGC CAG TGT GAA TGC AGA CCT AAA 383
Gly Glu Met Ser Leu Glu Glu His Ser Gln Cys Glu Cys Arg Pro Lys
115 120 125

AAA AAG GAC AGT GCT GTG AAG CCA GAT AGG TGC CGG AAG CTG CGA AGG 431
Lys Lys Asp Ser Ala Val Lys Pro Asp Arg Cys Arg Lys Leu Arg Arg
130 135 140

TGACACATGG CTTTTCAGAC TCAGCAGGGT GACTTGCCTC AGAGGCTATA TCCCAGTGGG 491

GGAACAAAGG GGAGCCTGGT AAAAAACAGC CAAGCCCCCA AGACCTCAGC CCAGGCAGAA 551

GCTGCTCTAG GACCTGGGCC TCTCAGAGGG CTCTTCTGCC ATCCCTTGTC TCCCTGAGGC 611

CATCATCAA CAGGACAGAG TTGGAAGAGG AGACTGGGAG GCAGCAAGAG GGGTCACATA 671

CCAGCTCAGG GGAGAATGGA GTACTGTCTC AGTTTCTAAC CACTCTGTGC AAGTAAGCAT 731

CTTACAAC TGCTCTTCCTC CCCTCACTAA GAAGACCCAA ACCTCTGCAT AATGGGATTT 791

GGGCTTTGGT ACAAGAACTG TGACCCCCAA CCCTGATAAA AGAGATGGAA GGAAAAAAAA 851

AAAAAAA 858

- 46 -

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 143 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met	Ser	Pro	Leu	Leu	Arg	Arg	Leu	Leu	Leu	Ala	Ala	Leu	Leu	Gln	Leu	1	5	10	15
Ala	Pro	Ala	Gln	Ala	Pro	Val	Ser	Gln	Pro	Asp	Ala	Pro	Gly	His	Gln	20	25	30	
Arg	Lys	Val	Val	Ser	Trp	Ile	Asp	Val	Tyr	Thr	Arg	Ala	Thr	Cys	Gln	35	40	45	
Pro	Arg	Glu	Val	Val	Val	Pro	Leu	Thr	Val	Glu	Leu	Met	Gly	Thr	Val	50	55	60	
Ala	Lys	Gln	Leu	Val	Pro	Ser	Cys	Val	Thr	Val	Gln	Arg	Cys	Gly	Gly	65	70	75	80
Cys	Cys	Pro	Asp	Asp	Gly	Leu	Glu	Cys	Val	Pro	Thr	Gly	Gln	His	Gln	85	90	95	
Val	Arg	Met	Gln	Ile	Leu	Met	Ile	Arg	Tyr	Pro	Ser	Ser	Gln	Leu	Gly	100	105	110	
Glu	Met	Ser	Leu	Glu	Glu	His	Ser	Gln	Cys	Glu	Cys	Arg	Pro	Lys	Lys	115	120	125	
Lys	Asp	Ser	Ala	Val	Lys	Pro	Asp	Arg	Cys	Arg	Lys	Leu	Arg	Arg	130	135	140		

- 47 -

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 910 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 3..305

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

CC ATG AGC CCT CTG CTC CGC CGC CTG CTG CTC GCC GCA CTC CTG CAG      47
Met Ser Pro Leu Leu Arg Arg Leu Leu Leu Ala Ala Leu Leu Gln
   1               5               10              15

CTG GCC CCC GCC CAG GCC CCT GTC TCC CAG CCT GAT GCC CCT GGC CAC      95
Leu Ala Pro Ala Gln Ala Pro Val Ser Gln Pro Asp Ala Pro Gly His
          20               25              30

CAG AGG AAA GTG GTG TCA TGG ATA GAT GTG TAT ACT CGC GCT ACC TGC      143
Gln Arg Lys Val Val Ser Trp Ile Asp Val Tyr Thr Arg Ala Thr Cys
          35               40              45

CAG CCC CGG GAG GTG GTG GTG CCC TTG ACT GTG GAG CTC ATG GGC ACC      191
Gln Pro Arg Glu Val Val Val Pro Leu Thr Val Glu Leu Met Gly Thr
          50               55              60

GTG GCC AAA CAG CTG GTG CCC AGC TGC GTG ACT GTG CAG CGC TGT GGT      239
Val Ala Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly
          65               70              75

GGC TGC TGC CCT GAC GAT GGC CTG GAG TGT GTG CCC ACT GGG CAG CAC      287
Gly Cys Cys Pro Asp Asp Gly Leu Glu Cys Val Pro Thr Gly Gln His
          80               85              90              95

CAA GTC CGG ATG CAG ACC TAAAAAAAG GACAGTGCTG TGAAGCCAGA      335
Gln Val Arg Met Gln Thr
          100

CAGGGCTGCC ACTCCCCACC ACCGTCCCCA GCCCGTTCT GTTCGGGCT GGGACTCTGC      395

CCCCGGAGCA CCCTCCCCAG CTGACATCAC CCATCCCACT CCAGCCCCAG GCCCCTCTGC      455

CCACGCTGCA CCCAGCACCA CCAGCGCCCT GACCCCGGA CCTGCCGCTG CCGCTGCCGA      515

CGCCGCAGCT TCCTCCGTTG CCAAGGGCGG GGCTTAGAGC TCAACCCAGA CACCTGCAGG      575

TGCCGGAAGC TGCGAAGGTG ACACATGGCT TTTCAGACTC AGCAGGGTGA CTTGCCTCAG      635

AGGCTATATC CAGTGGGGA ACAAAGAGGA GCCTGGTAAA AAACAGCCAA GCCCCCAAGA      695

```

- 48 -

```

CCTCAGCCCA GGCAGAAGCT GCTCTAGGAC CTGGGCCTCT CAGAGGGCTC TTCTGCCATC   755
CCTTGTCTCC CTGAGGCCAT CATCAAACAG GACAGAGTTG GAAGAGGAGA CTGGGAGGCA   815
GCAAGAGGGG TCACATACCA GCTCAGGGGA GAATGGAGTA CTGTCTCAGT TTCTAACCAC   875
TCTGTGCAAG TAAGCATCTT ACAACTGGCT CTTCC                                910

```

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 101 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

Met Ser Pro Leu Leu Arg Arg Leu Leu Leu Ala Ala Leu Leu Gln Leu
 1             5             10             15
Ala Pro Ala Gln Ala Pro Val Ser Gln Pro Asp Ala Pro Gly His Gln
      20             25             30
Arg Lys Val Val Ser Trp Ile Asp Val Tyr Thr Arg Ala Thr Cys Gln
      35             40             45
Pro Arg Glu Val Val Val Pro Leu Thr Val Glu Leu Met Gly Thr Val
      50             55             60
Ala Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly Gly
      65             70             75             80
Cys Cys Pro Asp Asp Gly Leu Glu Cys Val Pro Thr Gly Gln His Gln
      85             90             95
Val Arg Met Gln Thr
      100

```

- 49 -

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ACCACCACCT CCCTGGGCTG GCATGTGGCA CGTGCATAAA CG

42

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AGTTGTTTGA CCACATTGCC CATGAGTTCC ATGCTCAGAG GC

42

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 38 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GATCCTGGGG CTGGAGTGGG ATGGATGATG TCAGCTGG

38

- 50 -

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 40 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GCGGGCAGAG GATCCTGGGG CTGTCTGGCC TCACAGCACT

40

- 51 -

CLAIMS:

1. A biologically isolated proteinaceous molecule having the following characteristics:

- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one property in common with vascular endothelial growth factor (VEGF).

2. A proteinaceous molecule according to claim 1 wherein the molecule exhibits at least one of the following properties:

- (i) an ability to induce vascular endothelial cells;
- (ii) an ability to interact with *flt-1/flk-1* family of receptors; and/or
- (iii) an ability to induce cell migration, cell survival and/or an increase in intracellular levels of alkaline phosphatase.

3. A proteinaceous molecule according to claim 1 or 2 wherein said molecule has the capacity to induce astroglial proliferation.

4. A proteinaceous molecule according to claim 1 wherein said molecule is of human origin.

5. A proteinaceous molecule according to claim 1 wherein said molecule is of non-human origin.

6. A proteinaceous molecule according to claim 5 wherein said molecule is of livestock animal, companion animal, laboratory test animal, avian, fish or reptilian origin.

7. A proteinaceous molecule according to claim 5 wherein said molecule is encoded by a gene located at chromosome 11q13.

- 52 -

8. A proteinaceous molecule according to claim 1 wherein the percentage similarity to SEQ ID NO:2 is at least about 30%.
9. A proteinaceous molecule according to claim 1 wherein the percentage similarity to SEQ ID NO:2 is at least about 40%.
10. A proteinaceous molecule according to claim 1 wherein the percentage similarity to SEQ ID NO:2 is at least about 60-70%.
11. A proteinaceous molecule according to claim 1 comprising a sequence of amino acids as set forth in SEQ ID NO:4 or a part, fragment, derivative or analogue thereof.
12. A proteinaceous molecule according to claim 1 comprising an amino acid sequence substantially set forth in SEQ ID NO:6 or a part, fragment, derivative or analogue thereof.
13. A proteinaceous molecule according to claim 1 comprising an amino acid sequence substantially set forth in SEQ ID NO:8 or a part, fragment, derivative or analogue thereof.
14. A proteinaceous molecule according to claim 1 comprising an amino acid sequence substantially set forth in SEQ ID NO:10 or a part, fragment, derivative or analogue thereof.
15. A recombinant molecule having the following characteristics:
 - (i) an amino acid sequence substantially as set forth in SEQ ID NO:4 or having at least about 15% similarity to but at least about 5% dissimilarity to the amino acid sequence set forth in SEQ ID NO:2;
 - (ii) exhibits at least one biological property in common with VEGF.

16. A recombinant molecule having the following characteristics:
 - (i) an amino acid sequence substantially as set forth in SEQ ID NO:6 or having at least about 15% similarity to but at least about 5% dissimilarity to the amino acid sequence set forth in SEQ ID NO:2;
 - (ii) exhibits at least one biological property in common with VEGF.
17. A recombinant molecule having the following characteristics:
 - (i) an amino acid sequence substantially as set forth in SEQ ID NO:8 or having at least about 15% similarity to but at least about 5% dissimilarity to the amino acid sequence set forth in SEQ ID NO:2;
 - (ii) exhibits at least one biological property in common with VEGF.
18. A recombinant molecule having the following characteristics:
 - (i) an amino acid sequence substantially as set forth in SEQ ID NO:10 or having at least about 15% similarity to but at least about 5% dissimilarity to the amino acid sequence set forth in SEQ ID NO:2;
 - (ii) exhibits at least one biological property in common with VEGF.
19. A recombinant molecule according to claim 15 or 16 or 17 or 18 having at least one of the following properties:
 - (a) an ability to induce vascular endothelial cells;
 - (b) an ability to interact with *flt1/flki* family of receptors;
 - (c) an ability to induce cell migration, cell survival and/or increase intracellular levels of alkaline phosphatase.
20. A recombinant molecule according to claim 15 or 16 or 17 or 18 having the capacity to induce astroglial proliferation.

- 54 -

21. A recombinant molecule according to claim 20 wherein the molecule comprises an amino acid sequence substantially as set forth in SEQ ID NO:6.
22. A peptide fragment corresponding to a portion of the amino acid sequence set forth in SEQ ID NO:4 or a derivative or chemical equivalent thereof.
23. A peptide fragment according to claim 22 having the sequence set forth in SEQ ID NO:6 or a chemical equivalent thereof.
24. A peptide fragment according to claim 22 having the sequence set forth in SEQ ID NO:8 or a chemical equivalent thereof.
25. A peptide fragment according to claim 22 having the sequence set forth in SEQ ID NO:10 or a chemical equivalent thereof.
26. A nucleic acid molecule comprising a sequence of nucleotides or complementary to a sequence encoding a proteinaceous molecule having the following characteristics:
 - (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the sequence set forth in SEQ ID NO:2;
 - (ii) exhibits at least one property in common with vascular endothelial growth factor (VEGF).
27. A nucleic acid molecule according to claim 26 wherein the proteinaceous molecule exhibits at least one of the following properties:
 - (i) an ability to induce vascular endothelial cells;
 - (ii) an ability to interact with *flt-1/flk-1* family of receptors; and/or
 - (iii) an ability to induce cell migration, cell survival and/or an increase in intracellular levels of alkaline phosphatase.
28. A nucleic acid molecule according to claim 27 wherein the proteinaceous molecule has the capacity to induce astroglial proliferation.

- 55 -

29. A nucleic acid molecule according to claim 28 wherein said molecule encodes an amino acid sequence substantially as set forth in SEQ ID NO:6.
30. A nucleic acid molecule according to claim 1 wherein said molecule is of human origin.
31. A nucleic acid molecule according to claim 1 wherein the percentage similarity to SEQ ID NO:2 is at least about 30%.
32. A nucleic acid molecule according to claim 26 comprising a nucleotide sequence substantially as set forth in SEQ ID NO:3 or having at least 15% similarity thereto or capable of hybridising under low stringency conditions to a reverse complement of the nucleotide sequence as set forth in SEQ ID NO:3 provided that the nucleotide sequence has at least 15% similarity but at least 30% dissimilarity to the nucleotide sequence set forth in SEQ ID NO:3.
33. A nucleic acid molecule according to claim 26 encoding a murine homologue of human VEGF and comprising a nucleotide sequence substantially as set forth in Figure 9.
34. A pharmaceutical composition comprising a proteinaceous molecule according to claim 1 or 2 or 3 or 11 and one or more pharmaceutically acceptable carriers and/or diluents.
35. A method for preparing a recombinant molecule having the following characteristics:
- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the sequence set forth in SEQ ID NO:2;
 - (ii) exhibits at least one property in common with vascular endothelial growth factor (VEGF),
- said method comprising expressing a nucleic acid molecule encoding said recombinant

- 56 -

molecule by a suitable host grown under conditions effective to synthesise said recombinant molecule and then isolating said molecule.

36. A method according to claim 35 wherein the nucleic acid molecule comprises a sequence of nucleotides as set forth in SEQ ID NO:3 or having at least 15% similarity thereto or is capable of hybridising under low stringency conditions to a reverse complement of the nucleotide sequence as set forth in SEQ ID NO:3 provided that the nucleotide sequence has at least 15% similarity but at least 30% dissimilarity to the nucleotide sequence set forth in SEQ ID NO:3.

37. A method of inducing astroglial proliferation in a mammal, said method comprising administering to said mammal an effective amount of a recombinant proteinaceous molecule having the characteristics:

- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one property in common with vascular endothelial growth factor (VEGF),

said administration being for a time and under conditions sufficient to induce astroglial proliferation.

38. A method according to claim 37 wherein the recombinant proteinaceous molecule comprises an amino acid sequence substantially as set forth in SEQ ID NO:3 or is a derivative thereof.

39. A method according to claim 37 wherein the recombinant proteinaceous molecule comprises an amino acid sequence substantially as set forth in SEQ ID NO:6 or is a derivative thereof.

40. A method of promoting neuronal survival and/or proliferation in a mammal, said method comprising administering to said mammal an effective amount of a recombinant proteinaceous molecule having the characteristics:

- 57 -

- (i) comprises an amino acid sequence having at least about 15% similarity but at least about 5% dissimilarity to the sequence set forth in SEQ ID NO:2;
- (ii) exhibits at least one property in common with vascular endothelial growth factor (VEGF),

said administration being for a time and under conditions sufficient to induce astroglial proliferation.

41. A method according to claim 40 wherein the recombinant proteinaceous molecule comprises an amino acid sequence substantially as set forth in SEQ ID NO:3 or is a derivative thereof.

41. A method according to claim 40 wherein the recombinant proteinaceous molecule comprises an amino acid sequence substantially as set forth in SEQ ID NO:6 or is a derivative thereof.

1/52

<p>2/52</p> <p><i>Fig.1(i)</i></p>	<p>3/52</p> <p><i>Fig.1(ii)</i></p>
<p>4/52</p> <p><i>Fig.1(iii)</i></p>	<p>5/52</p> <p><i>Fig.1(iv)</i></p>

2/52

1	TCGGCCTCC GAAACC ATG AAC TTT CTG	
	Met Asn Phe Leu	1
50	CTT GCC TTG CTG CTC TAC CTC CAC	
	Leu Ala Leu Leu Leu Tyr Leu His	15
98	CCC ATG GCA GAA GGA GGA GGG CAG	
	Pro Met Ala Glu Gly Gly Gly Gln	30 35
146	ATG GAT GTC TAT CAG CGC AGC TAC	
	Met Asp Val Tyr Gln Arg Ser Tyr	45 50
194	GAC ATC TTC CAG GAG TAC CCT GAT	
	Asp Ile Phe Gln Glu Tyr Pro Asp	60 65
242	TCC TGT GTG CCC CTG ATG CGA TGC	
	Ser Cys Val Pro Leu Met Arg Cys	80
290	CTC GAG TGT GTG CCC ACT GAG GAG	
	Leu Glu Cys Val Pro Thr Glu Glu	95
338	CGG ATC AAA CCT CAC CAA GGC CAG	
	Arg Ily Lys Pro His Gln Gly Gln	110 115

Fig.1(i)

3/52

CTG TCT TGG GTG CAT TGG AGC Leu Ser Trp Val His Trp Ser 5 10	49
CAT GCC AAG TGG TCC CAG GCT GCA His Ala Lys Trp Ser Gln Ala Ala 20 25	97
AAT CAT CAC GAA GTG GTG AAG TTC Asn His His Glu Val Val Lys Phe 40	145
TGC CAT CCA ATC GAG ACC CTG GTG Cys His Pro Ile Glu Thr Leu Val 55	193
GAG ATC GAG TAC ATC TTC AAG CCA Glu Ile Glu Tyr Ile Phe Lys Pro 70 75	241
GGG GGC TGC TGC AAT GAC GAG GGC Gly Gly Cys Cys Asn Asp Glu Gly 85 90	289
TCC AAC ATC ACC ATG CAG ATT ATG Ser Asn Ile Thr Met Gln Ile Met 100 105	337
CAC ATA GGA GAG ATG AGC TTC CTA His Ile Gly Glu Met Ser Phe Leu 120	385

Fig.1(iii)

4/52

386	CAG CAC AAC AAA TGT GAA TGC AGA
	Gln His Asn Lys Cys Glu Cys Arg
	125 130
434	GAA AAT CCC TGT GGG CCT TGC TCA
	Glu Asn Pro Cys Gly Pro Cys Ser
	140 145
482	CAA GAT CCG CAG ACG TGT AAA TGT
	Gln Asp Pro Gln Thr Cys Lys Cys
	160
530	TGC AAG GCG AGG CAG CTT GAG TTA
	Cys Lys Ala Arg Gln Leu Glu Leu
	175
578	AAG CCG AGG CGG TGAGCCGGGC AGGAG
	Lys Pro Arg Arg
	190
630	GAACCAGATC TCTCACCAGG

Fig.1(iii)

5/52

CCA	AAG	AAA	GAT	AGA	GCA	AGA	CAA	433
Pro	Lys	Lys	Asp	Arg	Ala	Arg	Gln	
			135					

GAG	CGG	AGA	AAG	CAT	TTG	TTT	GTA	481
Glu	Arg	Arg	Lys	His	Leu	Phe	Val	
		150					155	

TCC	TGC	AAA	AAC	ACA	GAC	TCG	CGT	529
Ser	Cys	Lys	Asn	Thr	Asp	Ser	Arg	
	165					170		

AAC	GAA	CGT	ACT	TGC	AGA	TGT	GAC	577
Asn	Glu	Arg	Thr	Cys	Arg	Cys	Asp	
180					185			

GAAGG	AGCCTCCCTC	AGCGTTTCGG	629
-------	------------	------------	-----

649

Fig.1(iv)

6/52

7/52 <i>Fig.2(i)</i>	8/52 <i>Fig.2(ii)</i>
9/52 <i>Fig 2(iii)</i>	10/52 <i>Fig 2(iv)</i>
11/52 <i>Fig 2(v)</i>	12/52 <i>Fig 2(vi)</i>

7/52

1	CC ATG AGC CCT CTG CTC CGC CGC	
	Met Ser Pro Leu Leu Arg Arg	
	1 5	
48	CTG GCC CCC GCC CAG GCC CCT GTC	
	Leu Ala Pro Ala Gln Ala Pro Val	
	20	
96	CAG AGG AAA GTG GTG TCA TGG ATA	
	Gln Arg Lys Val Val Ser Trp Ile	
	35	
144	CAG CCC CGG GAG GTG GTG GTG CCC	
	Gln Pro Arg Glu Val Val Val Pro	
	50 55	
192	GTG GCC AAA CAG CTG GTG CCC AGC	
	Val Ala Lys Gln Leu Val Pro Ser	
	65 70	
240	GGC TGC TGC CCT GAC GAT GGC CTG	
	Gly Cys Cys Pro Asp Asp Gly Leu	
	80 85	
288	CAA GTC CGG ATG CAG ATC CTC ATG	
	Gln Val Arg Met Gln Ile Leu Met	
	100	
336	GGG GAG ATG TCC CTG GAA GAA CAC	
	Gly Glu Met Ser Leu Glu Glu His	
	115	

Fig.2(i)

8/52

CTG CTG CTC GCC GCA CTC CTG CAG Leu Leu Leu Ala Ala Leu Leu Gln 10 15	47
TCC CAG CCT GAT GCC CCT GGC CAC Ser Gln Pro Asp Ala Pro Gly His 25 30	95
GAT GTG TAT ACT CGC GCT ACC TGC Asp Val Tyr Thr Arg Ala Thr Cys 40 45	143
TTG ACT GTG GAG CTC ATG GGC ACC Leu Thr Val Glu Leu Met Gly Thr 60	191
TGC GTG ACT GTG CAG CGC TGT GGT Cys Val Thr Val Gln Arg Cys Gly 75	239
GAG TGT GTG CCC ACT GGG CAG CAC Glu Cys Val Pro Thr Gly Gln His 90 95	287
ATC CGG TAC CCG AGC AGT CAG CTG Ile Arg Tyr Pro Ser Ser Gln Leu 105 110	335
AGC CAG TGT GAA TGC AGA CCT AAA Ser Gln Cys Glu Cys Arg Pro Lys 120 125	383

Fig. 2(ii)

9/52

384	AAA	AAG	GAC	AGT	GCT	GTG	AAG	CCA
	Lys	Lys	Asp	Ser	Ala	Val	Lys	Pro
			130					135
432	CGT	CCC	CAG	CCC	CGT	TCT	GTT	CCG
	Arg	Pro	Gln	Pro	Arg	Ser	Val	Pro
		145					150	
480	CCC	TCC	CCA	GCT	GAC	ATC	ACC	CAT
	Pro	Ser	Pro	Ala	Asp	Ile	Thr	His
	160					165		
528	GCC	CAC	GCT	GCA	CCC	AGC	ACC	ACC
	Ala	His	Ala	Ala	Pro	Ser	Thr	Thr
				180				
576	GCT	GCC	GCT	GCC	GAC	GCC	GCA	GCT
	Ala	Ala	Ala	Ala	Asp	Ala	Ala	Ala
			195					

Fig. 2(iii)

10/52

GAC	AGG	GCT	GCC	ACT	CCC	CAC	CAC	431
Asp	Arg	Ala	Ala	Thr	Pro	His	His	
				140				
GGC	TGG	GAC	TCT	GCC	CCC	GGA	GCA	479
Gly	Trp	Asp	Ser	Ala	Pro	Gly	Ala	
				155				
CCC	ACT	CCA	GCC	CCA	GGC	CCC	TCT	527
Pro	Thr	Pro	Ala	Pro	Gly	Pro	Ser	
				170				175
AGC	GCC	CTG	ACC	CCC	GGA	CCT	GCC	575
Ser	Ala	Leu	Thr	Pro	Gly	Pro	Ala	
				185				190
TCC	TCC	GTT	GCC	AAG	GGC	GGG	GCT	T 624
Ser	Ser	Val	Ala	Lys	Gly	Gly	Ala	
200					205			

Fig.2(iv)

11/52

625	AGAGCTCAAC	CCAGACACCT	GCAGGTGCCG
685	GACTCAGCAG	GGTGACTTGC	CTCAGAGGCT
745	GGTAAAAAAC	AGCCAAGCCC	CCAAGACCTC
805	GCCTCTCAGA	GGGCTCTTCT	GCCATCCCTT
865	GAGTTGGAAG	AGGAGACTGG	GAGGCAGCAA
825	GGAGTACTGT	CTCAGTTTCT	AACCACTCTG
985	CTCCCCTCAC	TAAGAAGACC	CAAACCTCTG
1045	CTGTGACCCC	CAACCCTGAT	AAAAGAGATG

Fig.2(v)

12/52

GAAGCTGCGA	AGGTGACACA	TGGCTTTTCA	684
ATATCCCAGT	GGGGGAACAA	AGGGGAGCCT	744
AGCCCAGGCA	GAAGCTGCTC	TAGGACCTGG	804
GTCTCCCTGA	GGCCATCATC	AAACAGGACA	864
GAGGGGTCAC	ATACCAGCTC	AGGGGAGAAT	924
TGCAAGTAAG	CATCTTACAA	CTGGCTCTTC	984
CATAATGGGA	TTTGGGCTTT	GGTACAAGAA	1044
GAAGGAAAAA	AAAAAAAAAA		1094

Fig.2(vi)

13/52

14/52	15/52
Fig. 3(i)	Fig. 3(ii)

14/52

>VEGF_HUMAN VEGF_HUMAN VASCULAR ENDOTHELIAL
(VASCULAR 215 AA.
LENGTH = 215

SCORE = 181 (92.4 BITS), EXPECT = $6.4e-20$,
IDENTITIES = 33/75 (44%), POSITIVES = 48/75

QUERY: 31 HQRKVVSVIDVYTRATCQPREVVVPLTVEL
+++ VV +DVY R+ C+P E +V + E
SBJCT: 36 NHHEVVVKFMDVYQRSYCHPIETLVDIFQEY

QUERY: 91 PTGQHQVRMQILMIR 105
PT + + MQI+ I+
SBJCT: 96 PTEESNITMQIMRIK 110

SCORE = 76 (38.8 BITS), EXPECT = 0.0011,
IDENTITIES = 12/19 (63%), POSITIVES = 16/19

QUERY: 110 QLGEMSLEEHSQCECRPKK 128
++GEMS +H+ CECRPKK
SBJCT: 116 HIGEMSFLQHNKCECRPKK 134

SCORE = 72 (36.8 BITS), EXPECT = 0.0046,
IDENTITIES = 14/21 (66%), POSITIVES = 15/21

QUERY: 202 RCQGRGLELNPDTTCRCRKLRR 222
RC +R LELN TCRC K RR
SBJCT: 195 RCKARQLELNERTCRC DKPRR 215

SCORE = 46 (23.5 BITS), EXPECT = 47.,
IDENTITIES = 6/10 (60%), POSITIVES = 9/10

QUERY: 187 DPRTCRCRCR 196
DP+TC+C C+
SBJCT: 181 DPQTCKCSCK 190

SUBSTITUTE SHEET (RULE 26)

Fig.3(i)

15/52

GROWTH FACTOR PRECURSOR (VEGF)

 $P = 6.4e-20$
(64%)

MGTVAKQLVPSCVTVQRCGGCCPDDGLECV 90
+ PSCV + RCGGCC D+GLECV
PDEIEYIFKPSCVPLMRCGGCCNDEGLECV 95

 $\text{POISSON } P(2) = 9.1e-12$
(84%) $\text{POISSON } P(3) = 3.6e-18$
(71%) $\text{POISSON } P(4) = 7.3e-10$
(90%)*Fig. 3(i)*

16/52

17/52 <i>Fig.4(i)</i>	18/52 <i>Fig.4(ii)</i>
19/52 <i>Fig.4(iii)</i>	20/52 <i>Fig.4(iv)</i>

17/52

Gap Weight:3.00 Average Match:1.000
 Length Weight:0.100 Average Mismatch:-0.900
 Quality:100.9 Length:739
 Ratio:0.175 Gaps:30
 Percent Percent
 Similarity:69.703 Identity:69.703

```

28   ATGAGCCCTCTGCTCCGCCGCCTGC
      |||| | ||||| | ||
17   ATGAACTTTCTGCT.....GTCT..

68   TGCAGCTGGCCCCCGCCCAGGCCCC
      ||| ||| || | ||| |||
57   TGCTGCTCTACCTCCACCATGCCAA

118  CACCAGAGGA.....
      |||||
106  AGAAGGAGGAGGGCAGAATCATCAC

140  GTGTATACTCGC.GCTACCTGCCAG
      || ||| ||| |||| |||||
152  GTCTATCAGCGCAGCTA.CTGCCAT

194  T....GA.....CTGTGGAGCTCAT
      | || ||| ||| ||
201  TCCAGGAGTACCCTGATGAGATCGA

235  CCCAGCTGCGTGACTGTGCAGCGCT
      || ||| ||| | || |||
239  CCATCCTGTGTGCCCCCTGATGCGAT

285  CCTGGAGTGTGTGCCCCACTGGGCAG
      ||||| ||||| ||||| |||
289  CCTGGAGTGTGTGCCCCACTGAGGAG
  
```

Fig.4(i)

18/52

TGCTCGCCGCACT.....CC	67
...TGGGTGCATTGGAGCCTTGCCT	56
TGTCTCCCAGCCTGATGCCCCTGGC	117
GTGGTCCCAGGCTGCA.CCCATGGC	105
.AAGTGGTG....TCATGGATAGAT	147
GAAGTGGTGAAGTTCATG....GAT	151
CCCCGGGAG...GTGGTGGTGCCCT	193
CCAATCGAGACCCTGGTGGACATCT	200
GGGCACCGTGGCCAAACAGCTGGTG	234
GTACATCTT...CAA.....G	238
GTGGTGGCTGCTGCCCTGACGATGG	284
GCGGGGGCTGCTGCAATGACGAGGG	288
CACCAAGTCCGGATGCAGAT.....	329
TCCAACATCACCATGCAGATTATGC	338

Fig.4(ii)
SUBSTITUTE SHEET (RULE 26)

19/52

330CCTCATGATCCGGTACC
339	GGATCAAACCTCA.....C
369	GTCCCTGGAAGAACACAGCCAGTGT
376	GAGCTTCCTACAGCACAAACAATGT
419	GTGCTGTGAAGCCAGACAGGGCTGC
423	G.....AGCAAGACAAG.....
469	CGTTCTGTTCCGGGCTGGGACTCTG
443	...TGTGGGCCTTGCTCAGA.....
519	CATCACCCATCCCCTCCAGCCCCA
468
569	GC.....ACCACCAGCGCCC
469	GCATTTGTTTGTACAA.....
609	TGCCGACGCCGCAGCTTCCTCCGTT
509	TG.CAAAAACACAGACTC..GCGTT
657	AACCCAGACACCTGCAGGTGCCGGA
554	AACGAACGTACTTGCAGATGTGACA

Fig.4(iii)

20/52

CGAGCAGTCAGC...TGGGGGAGAT	368
CAAG..GCCAGCACATAGGAGAGAT	375
GAATGCAGACCTAAAAAAAAAGGACA	418
GAATGCAGACC...AAAGAAAGATA	422
CACTCCCCACCACCGTCCCCAGCCC	468
.....AAAATCCC.....	442
CCCCCGGAGCACCTCCCCAGCTGA	518
...GCGGAGAA.....	467
GGCCCCTCTGCCCACGCTGCACCCA	568
.....A	468
TGACCCCCGGACCTGCCGCTGCCGC	608
.GATCCGCAGACGTGTAAATGTTCC	508
GCCAAGGGCGGGGC..TTAGAGCTC	656
GC..AAGGCGAGGCAGCTTGAGTTA	553
AGCTGCGAAGGTGA	695
AGCCGAGGCGGTGA	592

Fig.4(iv)

21/52

<i>22/52</i> <i>Fig.5(i)</i>	<i>23/52</i> <i>Fig.5(ii)</i>	<i>24/52</i> <i>Fig.5(iii)</i>
<i>25/52</i> <i>Fig.5(iv)</i>	<i>26/52</i> <i>Fig.5(v)</i>	<i>27/52</i> <i>Fig.5(vi)</i>

22/52

165SOMSQ.MSF.msf MSF:687

Type: D Tuesday, June 20, 1995

Check:3140

	1
VEGF165	ATGAACTTTCTGCTGTCTTGGGTG
SOM175	ATGAGCCCTCTGCTCCGCCGCCTG
SOM175-e6	ATGAGCCCTCTGCTCCGCCGCCTG
SOM175-e6&7	ATGAGCCCTCTGCTCCGCCGCCTG
SOM175-e4	ATGAGCCCTCTGCTCCGCCGCCTG
	81
VEGF165	CACCCATGGCAGAAGGAGGAGGGC
SOM175	TGCCCCTGGCCACCAGAGGAAAGT
SOM175-e6	TGCCCCTGGCCACCAGAGGAAAGT
SOM175-e6&7	TGCCCCTGGCCACCAGAGGAAAGT
SOM175-e4	TGCCCCTGGCCACCAGAGGAAAGT
	161
VEGF165	CCAATCGAGACCCTGGTGGACATC
SOM175	GTGGTGGTGGCCCTTGACTG.TGGA
SOM175-e6	GTGGTGGTGGCCCTTGACTG.TGGA
SOM175-e6&7	GTGGTGGTGGCCCTTGACTG.TGGA
SOM175-e4	GTGGTGGTGGCCCTTGACTG.TGGA
	241
VEGF165	GATGCGATGCGGGGGCTGCTGCAA
SOM175	GCAGCGCTGTGGTGGCTGCTGCCC
SOM175-e6	GCAGCGCTGTGGTGGCTGCTGCCC
SOM175-e6&7	GCAGCGCTGTGGTGGCTGCTGCCC
SOM175-e4	GCAGCGCTGTGGTGGCTGCTGCCC

Fig.5(i)

23/52

CATTGGAGCCTTGCCTTGCTGCTCTACC
CTGCTCGCCGCACTCCTGCAGCTGGCCC
CTGCTCGCCGCACTCCTGCAGCTGGCCC
CTGCTCGCCGCACTCCTGCAGCTGGCCC
CTGCTCGCCGCACTCCTGCAGCTGGCCC

AGAATCATCACGAAGTGGTGAAGTTCAT
GGTGTCATGGATAGATGTGTATACTCGC
GGTGTCATGGATAGATGTGTATACTCGC
GGTGTCATGGATAGATGTGTATACTCGC
GGTGTCATGGATAGATGTGTATACTCGC

TTCCAGGAGTACCCTGATGAGATCGAGT
GCTCATGGGCACCGTGGCCAAAC..AGC
GCTCATGGGCACCGTGGCCAAAC..AGC
GCTCATGGGCACCGTGGCCAAAC..AGC
GCTCATGGGCACCGTGGCCAAAC..AGC

TGACGAGGGCCTGGAGTGTGTGCCCCACT
TGACGATGGCCTGGAGTGTGTGCCCCACT
TGACGATGGCCTGGAGTGTGTGCCCCACT
TGACGATGGCCTGGAGTGTGTGCCCCACT
TGACGATGGCCTGGAGTGTGTGCCCCACT

Fig. 5(iii)

24/52

80

TCCACCATGCCAAGTGGTCCCAGGCTG .
CCGCCCAGGCCCCTGTCTCCCAGCCTGA
CCGCCCAGGCCCCTGTCTCCCAGCCTGA
CCGCCCAGGCCCCTGTCTCCCAGCCTGA
CCGCCCAGGCCCCTGTCTCCCAGCCTGA

160

GGATGTCTATCAGCGCAGCTACTGCCAT
G CTACCTGC . CAGCC . CCGGGAG
G CTACCTGC . CAGCC . CCGGGAG
G CTACCTGC . CAGCC . CCGGGAG
G CTACCTGC . CAGCC . CCGGGAG

240

ACATCTTCAAGCCATCCTGTGTGCCCCCT
TGGTGCCCAG CTGCGTGACTGT
TGGTGCCCAG CTGCGTGACTGT
TGGTGCCCAG CTGCGTGACTGT
TGGTGCCCAG CTGCGTGACTGT

320

GAGGAGTCCAACATCACCATGCAGATTA
GGGCAGCACCAAGTCCGGATGCAGATCC
GGGCAGCACCAAGTCCGGATGCAGATCC
GGGCAGCACCAAGTCCGGATGCAGATCC
GGGCAGCACCAAGTCCGGATGCAGA . . .

Fig.5(iii)

25/52

	321
VEGF165	TGCGGATCAAACCTCACCAAGGCC
SOM175	TCATGATCCGG...TACCCGAGCA
SOM175-e6	TCATGATCCGG...TACCCGAGCA
SOM175-e6&7	TCATGATCCGG...TACCCGAGCA
SOM175-e4
	401
VEGF165	AAGAAAGATAG.....AGCAA
SOM175	AAAAAGGACAGTGCTGTGAAGCCA
SOM175-e6	AAAAAGGACAGTGCTGTGAAGCCA
SOM175-e6&7	AAAAAGGACAGTGCTGTGAAGCCA
SOM175-e4	AAAAAGGACAGTGCTGTGAAGCCA
	481
VEGF165AAGCA.....
SOM175	CTCTGCCCCCGGAGCACCCCTCCCC
SOM175-e6
SOM175-e6&7
SOM175-e4	CTCTGCCCCCGGAGCACCCCTCCCC
	561
VEGF165	A.....GATCCGCA
SOM175	GCACCACCAGCGCCCTGACCCCCG
SOM175-E6	GCACCACCAGCGCCCTGACCCCCG
SOM175-e6&7
SOM175-e4	GCACCACCAGCGCCCTGACCCCCG
	641
VEGF165	TTGAGTTAAACGAACGTACTTGCA
SOM175	TAGAGCTCAACCCAGACACCTGCA
SOM175-e6	TAGAGCTCAACCCAGACACCTGCA
SOM175-e6&7
SOM175-e4	TAGAGCTCAACCCAGACACCTGCA

Fig.5(iv)

26/52

AGCACATAGGAGAGATGAGCTTCCTACA
GTCAGCTGGGGGAGATGTCCCTGGAAGA
GTCAGCTGGGGGAGATGTCCCTGGAAGA
GTCAGCTGGGGGAGATGTCCCTGGAAGA
.....

GACAAGAA....AATCCCTGTGG.....
GACAGGGCTGCCACTCCCCACCACCGTC
GATAG.....
GATAG.....
GACAGGGCTGCCACTCCCCACCACCGTC

.....
AGCTGACATCACCCATCCCCTCCAGCC
.....CC
.....
AGCTGACATCACCCATCCCCTCCAGCC

GACGTGTAAATGTTCTGCAAAAAC.AC
GACCTGCCGCTGCCGCTGCCGACGCCGC
GACCTGCCGCTGCCGCTGCCGACGCCGC
.....
GACCTGCCGCTGCCGCTGCCGACGCCGC

687

GATGTGACAAGCCGAGGCGGTGA
GGTGCCGGAAGCTGCGAAGGTGA
GGTGCCGGAAGCTGCGAAGGTGA
.GTGCCGGAAGCTGCGAAGGTGA
GGTGCCGGAAGCTGCGAAGGTGA

Fig.5(v)

27/52

400

GCACAACAAATGTGAATGCAGACC...A
ACACAGCCAGTGTGAATGCAGACCTAAA
ACACAGCCAGTGTGAATGCAGACCTAAA
ACACAGCCAGTGTGAATGCAGACCTAAA
.....CCTAAA

480

.....GCCTTGCTCAGAGCGGAGA
CCCAGCCCCGTTCTGTTCCGGGCTGGGA
.....
.....
CCCAGCCCCGTTCTGTTCCGGGCTGGGA

560

.....TTTGTT.....TGTAC..A
CCAGGCCCCCTCTGCCCACGCTGCACCCA
CCAGGCCCCCTCTGCCCACGCTGCACCCA
.....
CCAGGCCCCCTCTGCCCACGCTGCACCCA

640

AGACTCG..CGTTGCAAGGCGAGGCAGC
AGCTTCCTCCGTTGCCAAGGGCGGGGCT
AGCTTCCTCCGTTGCCAAGGGCGGGGCT
.....
AGCTTCCTCCGTTGCCAAGGGCGGGGCT

Fig.5(vi)

28/52

<i>Fig 6(i)</i> 29/52	<i>Fig 6(ii)</i> 30/52
<i>Fig 6(iii)</i> 31/52	

29/52

VEGF ₁₆₅	M	N	F	L	L	S	W	V	H	W	S	L	A	L	L	L	Y	L	H	H	A	K	W	S	Q	A	A	P
SOM175 _{Short}	M	S	P	L	L	R	R	L	L	.	.	L	A	A	L	L	Q	L	A	P	A	Q	A	P
VEGF ₁₆₅	I	F	Q	E	Y	P	D	E	I	E	Y	I	F	K	P	S	C	V	P	L	M	R	C	G	G	C	C	N
SOM175 _{Short}	L	T	V	E	L	M	G	T	V	A	K	Q	L	V	P	S	C	V	T	V	Q	R	C	G	G	C	C	P
VEGF ₁₆₅	F	L	Q	H	N	K	C	E	C	R	P	K	K	D	R	A
SOM175 _{Short}	L	E	E	H	S	Q	C	E	C	R	P	K	K	K	D	S	A	V	K	P	D	R	A	A	T	P	H	H
VEGF ₁₆₅	C	K	C	S	C	K	N	T	D	S	R	C	K	A	R	Q	L	E	L	N	E	R	T	C	R	C	D	K
SOM175 _{Short}	H	A	A	P	S	T	S	A	L	T	P	G	P	A	A	A	A	A	A	D	A	A	S	S	V	A	K	
OR...																												
VEGF ₁₆₅	M	N	F	L	L	S	W	V	H	W	S	L	A	L	L	L	Y	L	H	H	A	K	W	S	Q	A	A	P
SOM175 _{Long}	M	S	P	L	L	R	R	L	L	.	.	L	A	A	L	L	Q	L	A	P	A	Q	A	P
VEGF ₁₆₅	I	F	Q	E	Y	P	D	E	I	E	Y	I	F	K	P	S	C	V	P	L	M	R	C	G	G	C	C	N
SOM175 _{Long}	L	T	V	E	L	M	G	T	V	A	K	Q	L	V	P	S	C	V	T	V	Q	R	C	G	G	C	C	P
VEGF ₁₆₅	F	L	Q	H	N	K	C	E	C	R	P	K	K	D	R	A
SOM175 _{Long}	L	E	E	H	S	Q	C	E	C	R	P	K	K	K	D	S	A	V	K	P	D	R	A	A	T	P	H	H
VEGF ₁₆₅	G	P	C	S	E	R	R	K	H	L	F	V	Q	D	P	Q	T	C	K	C	S	C	K	N	T	D	S	.
SOM175 _{Long}	P	R	C	T	Q	H	H	Q	R	.	.	P	D	P	R	T	C	R	C	R	C	R	R	R	S	F	L	.

Fig.6(ii)

M A E G G G Q N H E . V V K F M D V Y Q R S Y C H P I E T L V D 60
 V S Q P D A P G H Q R K V V S W I D V Y Q R A T C C P P R E V V P 55
 D E G L E C V P T E E S N I T M Q I M R I K P H Q G Q H I G E M S 121
 D D G L E C V P T E G Q H Q V R M Q I L M I R . Y P S S Q L G E M S 115
 R Q E N P C G P C S E R R K H L F . V Q D P Q T 170
 R P Q P R S V P G W D S A P G A P S P A D I T H P T P A P G P S A 175
 P R R
 G G A 191
 207

30/52

M A E G G G Q N H E . V V K F M D V Y Q R S Y C H P I E T L V D 60
 V S Q P D A P G H Q R K V V S W I D V Y Q R A T C C P P R E V V P 55
 D E G L E C V P T E E S N I T M Q I M R I K P H Q G Q H I G E M S 121
 D D G L E C V P T E G Q H Q V R M Q I L M I R . Y P S S Q L G E M S 115
 R Q E N P R P Q P R S V P G W D S A P G A P S P A D I T H P T P A P G P L C 170
 R C K A R Q L E L N E R T C R C D K P R R C 177
 R C Q G R G L E L N P D T C R C R K L R R 191
 222

Fig. 6(ii)

31/52

Areas of 100% homology are boxed and conserved residues thought to be involved in homodimerisation are underlined. The VEGF sequence depicted includes the 26 amino acid leader sequence (removal of which gives rise to mature VEGF₁₆₅) giving a total length of 191 amino acids.

Homology of SOM175 to VEGF₁₆₅ is 27% (33%) at the protein level, however within this are blocks of 100% homology. In particular, many structural residues are conserved including those thought to be involved in homodimerisation of VEGF (by comparison with PDGF).

ie. Cysteine-47

Proline-70, Cysteine-72, Valine-74

Arginine-77, Cysteine-78, Glycine-80, Cysteines-81 & 82

Cysteine-89, Proline-91

Cysteines 122 & 124

Fig.6(iii)

32/52

SPLICE VARIANTS OF SOM175




Fig. 7

33/52




Fig. 8A

34/52

5'UTR.... ATGAGG	*Exon 1 (60bp)	GGCCAG gtacgtgagg
tctccacacag GCCCCT	Exon 2 (43bp)	GGAAAG aatacttaca
tctgctccca TGGTGT	Exon 3 (187bp)	ATGCAG gtccgagatg
ctgaatacag ATCCTC	Exon 4 (73bp)	ATGCAG gtgtcaggca
acttttcaag ACCTAA	Exon 5 (34bp)	AGACAG gtgagtcttt
ctcctccgta GGCTGC	Exon 6 (101bp)	CTCCAG cccaggccc
cccactccag CCCCAG	Exon 7 (109bp)	ACCCAG acacctgtag
ccctgctcag GTGCCG	*Exon 8 (22bp)	AGGTGA ...3'UTR

Fig.8B

35/52

<i>36/52</i> <i>Fig. 9(i)</i>	<i>37/52</i> <i>Fig. 9(ii)</i>
<i>38/52</i> <i>Fig. 9(iii)</i>	<i>39/52</i> <i>Fig. 9(iv)</i>

36/52

-163 gcacgagctcaggccgctcgctgcggcgctg
-103 gggggccgcggaggagccgccccctgcgcc
-43 ggcggtctctggctgacccccccccacaccg

16 CGTCGCCTGCTGCTTGTTGCACTGCTGCAG
R R L L L V A L L Q

76 TTTGATGGCCCCAGTCACCAGAAGAAAGTG
F D G P S H Q K K V

136 ACATGCCAGCCCAGGGAGGTGGTGGTGCCT
T C Q P R E V V V P

196 AAACAAGTAGTGCCCAGCTGTGTGACTGTG
K Q L V P S C V T V

256 GGCCTGGAATGTGTGCCCAGTGGGCAACAC
G L E C V P T G Q H

316 TACCCGAGCAGTCAGCTGGGGGAGATGTCC
Y P S S Q L G E M S

376 CCTAAAAAAAAAGGAGAGTGCTGTGAGGCCA
P K K K E S A V R P

436 CAGCCCCGCTCTGTTCCGGGCTGGGACTCT
Q P R S V P G W D S

Fig.9(i)

37/52

cg ttg cg ctg cct g cg ccc aggg ct cg gga	
cc g ccc cgg gt ccc cgg gt cc g cg ccc at gg	
cc ggg ct aggg ccc g <u>ATGAGCCCCCTGCTG</u>	
M S P L L	-17
CTGGCTCGCACCAGGCCCTGTGTCCCAG	
L A R T Q A P V S Q	4
GTGCCATGGATAGACGTTTATGCACGTGCC	
V P W I D V Y A R A	24
CTGAGCATGGA ACTCATGGGCAATGTGGTC	
L S M E L M G N V V	44
CAGCGCTGTGGTGGCTGCTGCCCTGACGAT	
Q R C G G C C P D D	64
CAAGTCCGAATGCAGATCCTCATGATCCAG	
Q V R M Q I L M I Q	84
CTGGGAGAACACAGCCAATGTGAATGCAGA	
L G E H S Q C E C R	104
GACAGGGTTGCCATACCCACCCACCGTCCC	
D R V A I P H H R P	124
<u>ACCCCGGGAGCACCTCCCCAGCTGACATC</u>	
T P G A P S P A D I	144

Fig.9(iii)

38/52

496 ATCCATCCCACTCCAGCCCCAGGATCCTCT
 I H P T P A P G S S
 S P R I L

556 CTGACCCCCGGACCTGCCGTTGCCGCTGTA
 L T P G P A V A A V
 P D P R T C R C R C

616 GGGGCTTAGAGCTCAACCCAGACACCTGTA
 G A *
 R G L E L N P D T C

676 ctttccagactccacgggcccggctgcttt
736 agcacaggcgtaacctcctcagtctgggag
796 gagctctctcgccatcttttatctcccaga
856 atgtctcacctcaggggccagggtactctc
916 ttctggctggctgtctcccctcactatgaa
976 gggttctgttatgataactgtgacacacac
1036 gacactaaaaaaaaaaaaaaaaaaaaaaaaa

Fig.9(iii)

39/52

GCCCCGCTTGCACCCAGCGCCGCCAACGCC										
A	R	L	A	P	S	A	A	N	A	164
C	P	P	C	T	Q	R	R	Q	R	130
GACGCCGCGCTTCCTCCATTGCCAAGGGC										
D	A	A	A	S	S	I	A	K	G	184
R	R	R	R	F	L	H	C	Q	G	150
↓										
GGTGCCGGAAGCCGCGAAAGTGAcaagctg										
										186
R	C	R	K	P	R	K	*			167
tatggccctgcttcacagggagaagagtgg										
gtcactgccccaggacctggaccttttaga										
gctgccatctaacaattgtcaaggaacctc										
tcacttaaccaccctgggtcaagtgagcatc										
aaccctaaacttctaccaataacgggattt										
acacactcacactctgataaaagagatgga										
aaaaaaaaaaaa										

Fig.9(iv)

40/52

41/52	42/52
Fig 10(i)	Fig 10(ii)

41/52

A

hVRF167	-21	MSPLLRRLLLAALLQLAPAQAP	↓
mVRF167	-21	MSPLLRRLLLVALLQLARTQAP	
hVRF167	30	EVVVPLTVELMGTVAKQLVPSC	
		:	
mVRF167	30	EVVVPLSMELMGNVVKQLVPSC	
hVRF167	80	ILMIRYPSSQLGEMSLEEHSQC	
		:	
mVRF167	80	ILMIQYPSSQLGEMSLGEHSQC	
hVRF167	130	RPDPRTCRCRCRRRSFLRCQGR	
		:	
mVRF167	130	RPDPRTCRCRCRRRFLHCQGR	

B

hVRF186	116	RAATPHHRPQPRSVPGWDSAPG
mVRF186	116	RVAIPHHRPQPRSVPGWDSTPG
hVRF186	166	TPGPAAAAADAAASSVAKGGA*
		:
mVRF186	166	TPGPAVAVDAAASSIAKGGA*

Fig.10(i)

42/52

VSQPDAPGHQRKVVSVIDVYTRATCQPR 29

||| |:|:|:| | | | | | | | | |

VSQFDGPSHQKKVVPWIDVYARATCQPR 29

VTVQRCGGCCPDDGLECVPTGQHQVRMQ 79

| | | | | | | | | | | | | | | | | | | | | |

VTVQRCGGCCPDDGLECVPTGQHQVRMQ 79

ECRPKKKDSAVKPDSPRPLCPRCTQHHQ 129

| | | | | |:| |:| | | | | | | | | |:|

ECRPKKKESAVRPDSPRILCPPCTQRRQ 129

GLELNPDTCRCLRR* 167

| | | | | | | | | | | | | | | |

GLELNPDTCRCLPRK* 167

APSPADITHPTPAPGPSAHAAPSTTSAL 165

| | | | | | | | | | | | | |:| | | |

APSPADIIHPTPAPGSSARLAPSAANAL 165

186

186

Fig. 10(ii)

43/52

44/52	45/52
Fig 11(i)	Fig 11(ii)

44/52

mVRF167	-21	MSPLLRRLLVALLQL	:: :
mVEGF188	-26	MNFLLSVHWTLLLLYLHH	
mVRF167	25	TCQPREVVVPLSMELMGNVV	: : ::
mVEGF188	24	YCRPIETLVDIFQEYPDEIE	
mVRF167	75	QVRMQILMIQYPSSQLGEM	: : :
mVEGF188	74	NITMQIMRIKPHQSQHIGEM	
mVRF167	119ILCPPC	:
mVEGF188	124	QKRKRKKSFRKSWSVHCEPC	
mVRF167	152	GLELNPDTCRCKPRK	:
mVEGF188	173	QLELNERTCRCDKPRR	

Fig.11(i)

45/52

AR.TQAPVSQFDGPSHQKKVVPWIDVYARA	24
: : : : : : ::	
AKWSQAAPTT.EGEQKSHEVIKFMDVYQRS	23
KQLVPSCVTVQRCGGCCPDDGLECVPTGQH	74
: : : : : : :	
YIFKPSCVPLMRCAGCCNDEALECVPTSES	73
SLGEHSQCECRPKKKESAVRPDSPR.....	118
: :	
SFLQHSRCECRPKKDRTKPEKKSVRGKGKG	123
TQRRQR...PDPRTCRCRCRRRFLHCQGR	151
: : : : : :	
SERRKHLFVQDPQTCKCSCKNTDS.RCKAR	172
	167
	188

Fig.11(ii)

46/52




Fig.12

47/52




Fig.13

48/52




Fig.14

49/52




Fig. 15

50/52




Fig. 16

51/52




Fig.17

SUBSTITUTE SHEET (RULE 26)

52/52

MOUSE ASTROGLIAL CELLS




Fig.18

MOUSE OLIGODENDROGLIAL CELLS




Fig.19

MOUSE FOREBRAIN NEURONS




Fig.20

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/AU 96/00094

A. CLASSIFICATION OF SUBJECT MATTER

Int Cl⁶: C12N 15/12; C07K 14/475; A61K 037/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

WPAT AND CHEM ABS

SEE DETAILS IN ELECTRONIC DATABASE BOX BELOW

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPM, JAPIO

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
DERWENT WPAT, USPM, JAPIO DATABASES; KEYWORDS: MVRF OR HVRF OR SOM 1: OR SOM X:
OR [VASOACTIVE () PERMEABILITY () FACTOR#] OR VEGF: OR VEGF OR VRF: OR VRF OR
[GROWTH () FACTOR (5N) (VASCULAR OR ENDOTHELI:)]
CHEMICAL ABSTRACTS DATABASE; KEYWORDS: AS ABOVE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AU 60798/90 (CALIFORNIA BIOTECHNOLOGY INC) published 21 February 1991	1-41
X	AU 56574/90 (GENENTECH, INC.) published 15 November 1990	1-41
P,X	AU 73941/94 (HUMAN GENOME SCIENCES, INC.) published 14 September 1995	1-41



Further documents are listed in the continuation of Box C



See patent family annex

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search
03 June 1996

Date of mailing of the international search report

5TH JUNE 1996.

Name and mailing address of the ISA/AU
AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION
PO BOX 200
WODEN ACT 2606
AUSTRALIA Facsimile No.: (06) 285 3929

Authorized officer

ARATI SARDANA

Telephone No.: (06) 283 2627

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/AU 96/00094

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5073492 (Chung-Ho Chen and Sumi C. Chen) published 17 December 1991	34
X	Biochemical and Biophysical Research Communications (1992), Vol 183, No. 3, pg 1167-1174 (Weindel K. et al.) "AIDS-ASSOCIATED KAPOSI'S SARCOMA CELLS IN CULTURE EXPRESS VASCULAR ENDOTHELIAL GROWTH FACTOR". See whole Article.	1-41
X	Biochemical and Biophysical Research Communications (1989), Vol 165, No. 3 pg 1198-1206 (Edmund Tischer et al.) "VASCULAR ENDOTHELIAL GROWTH FACTOR : A NEW MEMBER OF THE PLATELET-DERIVED GROWTH FACTOR GENE FAMILY". See whole Article.	1-41
X	Journal of Virology (1994), Vol 68, No. 1 pg 84-92 (David J. Lytle et al.) "HOMOLOGS OF VASCULAR ENDOTHELIAL GROWTH FACTOR ARE ENCODED BY THE POX VIRUS ORF VIRUS". See whole Article.	1-41
X	Methods in Enzymology (1991), vol 198, pg 391-405 (Ferrara Napoliana et al.) "PURIFICATION AND CLONING OF VASCULAR ENDOTHELIAL GROWTH FACTOR SECRETED BY PITUITARY FOLLICULOSTELLATE CELLS". See whole Article	1-41
X	The Journal of Biological Chemistry (1991), vol 266, No. 18 pg 11947-11954 (Edmund Tischer et al.) "THE HUMAN GENE FOR VASCULAR ENDOTHELIAL GROWTH FACTOR". See whole Article	1-41
P,X	Biochemical and Biophysical Research Communications (1996), Vol 220, No. 1 pg 147-52 (Lagercrantz J et al) "EXPRESSION OF THE VEGF-RELATED FACTOR GENE IN PRE- AND POSTNATAL MOUSE". See whole Article	33
P,X	Biochimica et Biophysica Acta (1995), Vol. 1260 No. 2 pg 235-9 (Sharma Hari S et al.) "NUCLEOTIDE SEQUENCE AND EXPRESSION OF THE PORCINE VASCULAR ENDOTHELIAL GROWTH FACTOR". See whole Article.	1-41
X	DEVELOPMENT (1992), Vol 114, pg 521-532 (Breier G et al.) "EXPRESSION OF VASCULAR ENDOTHELIAL GROWTH FACTOR DURING EMBRYONIC ANGIOGENESIS AND ENDOTHELIAL CELL DIFFERENTIATION". See whole Article	1-41
X	Molecular Endocrinology (1991), Vol 5 No. 12, pg 1806-1814 (Houck Keith A et al.) "THE VASCULAR ENDOTHELIAL GROWTH FACTOR FAMILY: IDENTIFICATION OF A FOURTH MOLECULAR SPECIES AND CHARACTERIZATION OF ALTERNATIVE SPLICING OF RNA". See whole Article	1-41

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/AU 96/00094

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
AU	56574/90	US IL WO	5332671 94128 9013649	CA JP	2054699 4505705	EP NZ	471754 233451
AU	60798/90	US JP	5194596 5501350	CA WO	2063810 9102058	EP US	484401 5219739
AU	73941/94	WO	9524473				
US	5073492	NONE					